### THE ZOOLOGICAL RECORD

VOLUME 96 SECTION 11 1959

Z 7991 Z87 V. 96

# TRILOBITA

COMPILED BY

J. T. TEMPLE, M.A., Ph.D.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE EIGHT SHILLINGS
1961



It will greatly help in the production of The Zoological Record and assist the Recorders of the individual sections, if authors will forward a copy of their paper or memoir to the Editor of the Zoological Record, The Zoological Society of London, Regent's Park, London, N.W.1. In the case of separately printed copies of articles so forwarded, the original pagination should be given.

All business correspondence concerning the Record should be addressed to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

### THE MUSEUMS JOURNAL

PUBLISHED BY THE MUSEUMS ASSOCIATION
22 Fitzroy Street, Fitzroy Square, London, W.1.

This journal, which is published monthly, contains articles, reviews and technical notes on every Museum aspect of zoology and other sciences. It is the only English journal which deals with the installation and preservation of exhibits, and which reviews Museum work in all parts of the world.

It is published monthly price 4s., by the Association and can be obtained from the above address. It is distributed free to all members of the Museums Association (Subscription £3 3s. 0d. per annum), of which full particulars are obtainable from the Secretary at the above address.

## **TRILOBITA**

COMPILED BY

J. T. TEMPLE, M.A., Ph.D.

fic

## II. TRILOBITA

COMPILED BY

J. T. TEMPLE, M.A., Ph.D.

#### CONTENTS

									PAGE
I.	TITLES	••••		****	****	****	****	****	2
II.	SUBJECT	INDEX	****	****	****	****	****	****	6
Ш.	SYSTEMA	TIC IND	EX	****	****	****			9
IV.	ERRATUN	£							32

#### I.—TITLES

Opinion 537. Designation under the Plenary Powers of type species in harmony with accustomed usage for the genera Encrinurus Emmrich, 1844, and Odontochile Hawle & Corda, 1847, and validation under the same Powers of the specific name punctatus Wahlenberg, 1821, as published in the combination Entomostracites punctatus (Class Trilobita). Opin. int. Comm. 2001. Nom. 20 4 1959: 43-56 pl. i.

Opinion 538. Designation under the Plenary Powers of a type species in harmony with accustomed usage for the nominal genus *Ptychopyge* Angelin, 1854 (Class Trilobita). Opin. int. Comm. zool. Nom. 20 5 1959: 59-64.

Anatol'eva, A. I. & Shelkovnikov, A. D. [On Middle Cambrian red beds of the south-western slope of Eastern Sayan.] C. R. Acad. Sci. U.R.S.S. (ns) 127 3 1959: 624-626 text-fig. 1 [In Russian] [Trilobita det. A. G. Sivov].

Badillo, L. Catálogo de especies fósiles del Museo del Instituto Geológico y Minero de España. 1.—Cambriano. Notas Inst. geol. Esp. 55 1959: 71–124 50 pls. map.

Balashov, Z. G. see Balashova, E. A.

Balashova, E. A. (1). [Middle and Upper Ordovician and Lower Silurian trilobites of Eastern Taimyr and their stratigraphical significance.] [Part 1]. Sborn. stat. paleont. biostrat. Inst. geol. Arktiki, Leningrad 14 1959: 17-47 pls. i-vi [In Russian].

Balashova, E. A. (2). [Middle and Upper Ordovician and Lower Silurian trilobites of Eastern Taimyr and their stratigraphical significance.] [Part 2]. Sborn. stat. paleont. biostrat. Inst. geol. Arktiki, Leningrad 15 1959: 27–55 [In Russian] [Part 3 of the text (Ibid. 17: 5-40), although dated 1959, seems to have appeared in 1960 and is not recorded here].

Balashova, E. A. & Balashov, Z. G. [On the stratigraphy of the Glauconitic and Orthoceratitic series of the Ordovician of the north-west of the Russian platform.] Ann. Leningr. Univ. No. 268 (Geol. Sci. No. 10) 1959: 127-154 table [In Russian].

Balk, C. Lochman see Lochman, C.

Berg, R. R. & Ross, R. J., Jnr. Trilobites from the Peerless and Manitou formations, Colorado. J. Paleont. 33 1 1959: 106-119 pls. xxi-xxii text-figs.

[ I

We Pro l-1

letz 195

I Pse Pro

gen

5 2

5-1

198

roc

lim

262

we

Spe

[ed

Ar

An G.

C. E.

C.

W

fro

J.

ta

Bodylevsky, V. I. [Little atlas of leading fossils.] 2nd ed. Leningrad & Moscow 1953: 239 pp. 86 pls. (drawings) [Trilobita pls. ii-v, viii, xiii, xiv] [In Russian].

Bordet, P., Cavet, P. & Pillet, J. Sur l'existence d'une faune d'âge silurien dans la région de Kathmandu (Himalaya du Népal). C. R. Acad. Sci., Paris 248 10 1959 : 1547—1549.

Borukaev, R. A. et al. [Resolutions of the Conference on the unification of the stratigraphical schemes of the Prepalaeozoic and Palaeozoic of Eastern Kazakhstan.] Acad. Sci. Kazakhstan S.S.R., Alma-Ata 1958: 39 pp. 6 tables [In Russian].

Bright, R. C. A paleoecologic and biometric study of the Middle Cambrian trilobite *Elrathia kingii* (Meek). J. Paleont. 33 1 1959: 83–98 pls. xvii–xviii text-figs. 1–8 tables 1–2.

Butcher, N. E. see Dearman, W. R.

Caldwell, W. G. E. The Lower Carboniferous rocks of the Carrick-on-Shannon syncline. Quart. J. geol. Soc. Lond. 115 2 1959: 163–186 pl. vi text-figs. 1–6.

Cavet, P. see Bordet, P.

Chang, W. T. New Trilobites from the Middle Cambrian of North China. Acta palaeont. sinica 7 3 1959: 193-214 [Chinese], 215-236 [English] pls. i-iv text-figs. 1-27.

**Colquhoun,** D. J. Stratigraphy and Palaeontology of the Nipissing—Deux Rivières Outliers. Proc. geol. Ass. Canada **10** 1958: 83–93 text-fig. 1 tables.

Coma, E. Suñer see Suñer Coma, E.

Dean, W. T. (1). The Stratigraphy of the Caradoc Series in the Cross Fell Inlier. Proc. Yorks, gool. Soc. 32 2 1959: 185-226 text-figs. 1-6.

Dean, W. T. (2). *Duftonia*, a new trilobite genus from the Ordovician of England and Wales. Palaeontology 2 1 1959: 143-149 pl. xix.

Dearman, W. R. & Butcher, N. E. The Geology of the Devonian and Carboniferous Rocks of the North-West Border of the Dartmoor Granite, Devonshire. Proo. Geol. Ass. 70 1 1959: 51-90 pl. ii text-figs. 1-14 [Trilobita det. R. Goldring].

Erben, H. K. Fortschritte der Paläontologie im letzten Jahrzehnt. Naturw. Rdsch., Stuttgart 12 4 1959: 119-124.

Erben, H. K. see Pillet, J.

Fernández de Villalta, J. see Suñer Coma, E.

Fritz, M. A. A Larval Stage of the Trilobite Pseudogygites latimarginatus (Hall) [Abstract]. Min. Proc. roy. Soc. Can. (3) 53 Appendix C 1959: 24.

Glaessner, M. F. & Parkin, L. W. [eds.] The geology of South Australia. J. geol. Soc. Australia 5 2 (for 1957) 1958: 1–163 11 pls. 5 maps 26 text-figs.

Goldring, R. see Dearman, W. R.

**Gómez de Llarena**, J. Rudolf Richter (7-XI-1881—5-I-1957) y Emma Richter (4-III-1888—15-XI-1956). Bol. Soc. esp. Hist. nat. [Geol.] **55** 1958: 17-25 portraits.

Hadding, A. The pre-quaternary sedimentary rocks of Sweden. VII. Cambrian and ordovician limestones. Lunds Univ. Årsskr. (nF) [2] 54 5 1958; 262 pp. 193 text-figs.

Hamada, T. Gotlandian Shelly Fauna from Southwest Japan (I). Coronocephalus kobayashii, a new Species from the Kuraoka District, Kyūshū. Japan. J. Geol. Geogr. 30 7 1959; 71-88 pl. vi text-figs. 1-2.

Harper, J. C. see Shackleton, R. M.

Harrington, H. J. et al. Trilobita, in R. C. Moore [ed.], Treatise on Invertebrate Paleontology Part O Arthropoda 1. Univ. Kansas Press & Geol. Soc. Amer. 1959: 038-0560 text-figs. 27-415 [Co-authors G. Henningsmeen, B. F. Howell, V. Jaanusson, C. Lochman-Balk, R. C. Moore, C. Poulsen, F. Rasetti, E. & R. Richter, H. Schmidt, K. Sdzuy, W. Struve, C. J. Stubblefield, R. Tripp, J. M. Weller, & H. B. Whittington].

Henderson, G. G. L. see North, F. K.

**Henningsmoen,** G. Rare Tremadocian Trilobites from Norway. Norsk geol. Tidsskr. **39** 2–3 1959: 153–174 pls. i–ii text-fig. 1.

Henningsmoen, G. see Harrington, H. J.

Holland, C. H. The Ludlovian and Downtonian rocks of the Knighton district, Radnorshire. Quart, J. geol. Soc. Lond. 114 4 (for 1958) 1959: 449-478 pl. xxi text-figs. 1-4 tables I-II.

Holland, C. H., Lawson, J. D. & Walmsley, V. G. A Revised Classification of the Ludlovian Succession at Ludlow. Nature, Lond. **184** 4692 1959; 1037–1039 table 1.

Howell, B. F. see Harrington, H. J.

Hu, Chung-Hung see Lochman, C.

Ivanova, E. A. [Development of the fauna of the Middle and Upper Carboniferous soa of the western part of the Moscow syncline in connection with its history. 3 Development of the fauna in connection with conditions of existence.] Trud. paleont. Inst. Acad. Sci. S.S.S.R. 69 1958: 1-303 pls. i-xxi frontispiece text-figs. 2-77 [In Russian].

Ivshin, N. K. [Eight new genera of trilobites from the Upper Cambrian of Central Kazakhstan.] Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955: 106–123 pl. i [In Russian with summary in Kazakh].

Jaanusson, V. see Harrington, H. J.

Jones, W. P. New species of phacopid trilobite [Abstract]. Bull. geol. Soc. Amer. 70 12 1959: 1727.

Kaljo, D. (1). Eesti geoloogilisest arenemisest vanaaegkonnas [The Geological Development of Estonia in the Palaeozoic]. Eesti Loodus 1959 4: 193-199 pls. xxiv-xxv text-figs. 1-4.

Kaljo, D. (2). Kivististest ja nende kogumisest [Fossils and How to Collect them]. Eesti Loodus 1959 4: 245-250 pls. xxix-xxx text-figs. 1-8.

Kauffman, M. E. Cambrian stratigraphy in the Garnet Range of western Montana [Abstract]. Bull. geol. Soc. Amer. 70 12 1959: 1780-1781.

Kegel, W. Nachruf für Rudolf Richter—Emma Richter. Z. dtsch. geol. Ges. 110 3 (for 1958) 1959: 637-642 portraits.

Khomentovsky, V. V. see Zhuravleva, I. T.

Kindle, C. H. & Whittington, H. B. Some stratigraphic problems of the Cow Head area in Western Newfoundland. Trans. N.Y. Acad. Sci. (2) 22 1 1959: 7–18 text-figs. 1–3.

Koroleva, M. N. [New genera of trilobites from the Middle and Upper Ordovician of northern Kazakhstan.] C. R. Acad. Sci. U.R.S.S. (ns) 124 6 1959: 1313-1316 pl. text-fig. 4 [In Russian].

Kravtsov, A. G. see Miroshnikov, L. D.

Lawson, J. D. see Holland, C. H.

Lazarenko, N. P. [Middle Cambrian Pagetides of the northern Siberian platform (Trilobites).] Sborn. stat. paleont. biostrat. Inst. geol. Arktiki, Leningrad 14 1959: 5-15 pl. i table I [In Russian].

Lesnikova, A. F. & Weber, V. N. [Class Trilobita, in M. E. Yanishevsky (ed.), Atlas of the leading forms of the fossil faunas of the USSR Volume II Silurian System.] VSEGEI, Moscow 1949: 270-315 pls. lxiv-lxxxv [In Russian: photostat only seen].

Lewowicki, S. Fauna wapieni klimeniowych z Dzikowca Kłodzkiego [Fauna of Clymenia limestones from Dzikowice near Kłodzko (Lower Silesia)]. Biul. Inst. Geol. [Bull. Serv. géol. Pologne] 146 1959: 73–112 [Polish text], 113–115 [Russian summary], 116–118 [English summary] pls. i–ii text-figs. 1–2.

Lochman, C. & Hu, Chung-Hung. A Ptychaspis faunule from the Bear River Range, southeastern Idaho. J. Paleont. 33 3 1959: 404-427 pls. lvii-lx text-fig.

Lochman-Balk, C. see Harrington, H. J.

Lu Yen-hao. Subdivision and correlation of the Ordovician rocks of South China. Geology Publishing House, Peking 1959: 1-105 [Chinese text], 106-113 [English summary] tables 1-16.

McKee, E. D. Cambrian History of the Grand Canyon Region Part I. Stratigraphy and Ecology of the Grand Canyon Cambrian. Publ. Carneg. Instn No. 563 1945: 5-168 pls. i-xv text-figs. 1-12 tables [Fossil lists det. C. E. Resser q. v.].

McLaren, D. J. The Role of Fossils in Defining Rock Units with Examples from the Devonian of Western and Arctic Canada. Amer. J. Sci. 257 10 1959: 734-751 text-figs. 1-8.

McLaren, D. J. see Thorsteinsson, R.

McWhae, J. R. H. et al. The stratigraphy of Western Australia. J. geol. Soc. Australia 4 2 (for 1956) 1958: 1-161 map 10 text-figs. 9 tables.

Maksimova, Z. A. & Organova, N. M. [The first discovery of remains of a Devonian fauna in Western Primor'e.] C. R. Acad. Sci. U.R.S.S. (ns) 128 3 1959: 594-595 text-fig. 1 [Trilobita det. Maksimova] [In Russian].

Männil, R. [M.] [On the stratigraphy of the Nabala horizon (F1a) of the Upper Ordovician of the Estonian SSR.] Geol. Inst. Uurimused [Trud. Inst. Geol. Acad. Sci. Est. SSR], Tallinn 2 1958: 3–15 figs. 1–2 [In Russian, with Estonian and German summaries pp. 15–17].

Miroshnikov, L. D., Kravtsov, A. G. & Shcheglova, O. S. [An outline of the stratigraphy of the Lower and Middle Palaeozoic of the north-western outskirts of the Siberian Platform.] C. R. Acad. Sci. U.R.S.S. (ns) 126 2 1959: 359–362 [Trilobita det. various authors] [In Russian].

Monseur, G. Sur un Trilobite déterminé comme Asteropyge punctata (Steininger). Ann. Soc. géol. Belg. 82 [Bull.] 1958: 89-96 pl.

Moore, R. C. see Harrington, H. J.

North, F. K. & Henderson, G. G. L. Summary of the Geology of the Southern Rocky Mountains of Canada. Alberta Soc. Petrol. Geol., 4th ann. Field Conf. Guidebook [1954]: 15-81 pls. v-xvi text-fig. 1 correlation chart.

Öpik, A. A. (1). Genal Caeca of Agnostids. Nature, Lond. 183 4677 1959: 1750-1751 text-fig. 1.

Öpik, A. A. (2). Tumblagooda sandstone trails and their age. Rep. Bur. Min. Res. Aust. No. 38 1959: 3–20 11 pls. [figs. 2–19] map.

Organova, N. M. see Maksimova, Z. A.

Orlowski, S. (1). On the Presence of Paradoxides ölandicus Beds in the Holy Cross Mountains. Bull. Acad. polon. Sci. [3] 5 7 1957: 769–772 pl.

Orlowski, S. (2). Paradoxidae from lower Middle Cambrian Strata in the Vicinity of Sandomierz (Central Poland). Bull. Acad. polon. Sci. [sci. chim., géol., géogr.] 7 6 1959: 441-446 pls. i-ii.

Orlowski, S. (3). Ellipsocephalidae from the Lower Beds of the Middle Cambrian in the Vicinity of Sandomierz (Central Poland). Bull. Acad. polon. Sci. [sci. chim., géol., géogr.] 7 7 1959: 515-520 pls. i-ii text-fig. 1.

Parkin, L. W. see Glaessner, M. F.

Paulus, B. Der mittlere Teil der Sötenicher Mulde (Devon, Eifel). I. Unterdevon und tiefes Eiflium. Senckenbergiana leth. 40 5–6 1959: 333–362 pls. i-ii text-figs. 1–6.

Petrakov, V. U. [New data on the stratigraphy of the Cambro-Ordovician deposits of the basin of the river Kureika.] C. R. Acad. Sci. U.R.S.S. (ns) 127 3 1959: 651-654 text-figs. 1-2 [In Russian].

**Pfeiffer,** H. Neue Beobachtungen und Funde aus dem Saalfelder Oberdevon. Geologie **8** 3 1959: 262–272 pls. i–iv table 1.

Pillet, J. (1). Contribution à l'étude du Dévonien Armoricain II.—La faune du Siegenien inférieur en Anjou. Bull. Soc. Étud. sci. Angers 87 année (1954-1957) 1958: 15-25.

Pillet, J. (2). Trilobites, in H. K. Erben et al., Note préliminaire sur la faune des calcaires de la Grange (M.-et-L.). Bull. Soc. Étud. sci. Angers (ns) 2 1959 : 102–103 [Observations also by H. K. Erben].

Pillet, J. see Bordet, P.

Pokrovskaya, N. V. [Trilobite fauna and stratigraphy of the Cambrian sediments of Tuva.] Trud. geol. Inst. [GIN] S.S.S.R. 27 1959: 1–200 pls. i–xi text-figs. 1–6 tables errata slip [In Russian].

Pokrovskaya, N. V. see Zhuravleva, Z. A.

Poulsen, C. see Harrington, H. J.

Prantl, F. Review of M. Šnajdr, Trilobiti českého středního kambria [recorded Z. R. 95 11]. Čas. nár. Mus. [Přír.] 127 2 1958: 234–235.

Rasetti, F. Trempeleauian trilobites from the Concocheague, Frederick, and Grove limestones of the central Appalachians. J. Paleont. 33 3 1959: 375–398 pls. li–lv text-figs. 1–2.

Rasetti, F. see Harrington, H. J.

Regnéll, G. The Pre-Quaternary Fossil-Bearing Systems Outside the High Mountains.... Geol. Fören. Stockh. Förh. 80 4 (for 1958) 1959 : 407–422.

Rehoř, F. & Rehořová, M. Trilobit Weberides mucronatus (M'Coy, 1844) v ostravském namuru [Trilobit Weberides mucronatus (M'Coy, 1844) im Ostrauer Namur]. Přírod. Čas. Sleský 20 3 1959: 254–271 pls. i–iv text-figs. 1–4 [German summary p. 272].

Řehořová, M. see Řehoř, F.

Repina, L. N. see Zhuravleva, I. T.

Repina, L. N. see Zhuravleva, Z. A.

Resser, C. E. Cambrian History of the Grand Canyon Region Part II. Cambrian Fossils of the Grand Canyon. Publ. Carneg. Instn No. 563 1945: 171–220 pls. xvi-xxvii. Begin

Dal

lone

[19

R

R

R

R

Upr

183.

R

No.
SCar
Ace

pol 524 opg An

gé

Car

u.

Do

30

K (n R

H te w H

F

Resser, C. E. see McKee, E. D.

Richter, E. see Harrington, H. J.

Richter, R. see Harrington, H. J.

Richter, R. & E. see Gómez de Llarena, J.

Richter, R. & E. see Kegel, W.

Robison, R. A. A Preliminary Correlation of the Upper Cambrian Faunas of Central and Western Utah [Abstract]. Proc. Utah Acad. Sci. 35 1958: 183.

Ross, R. J., Jnr. see Berg, R. R.

Rusconi, C. El Cámbrico medio de Mendoza según Poulsen. Rev. Mus. Hist. nat. Mendoza 12 1–4 1959 : 3–12.

Sampelayo, P. H. De la fauna gotlandiense: Dalmanites Batalleri, Samp.; Corrección del Phacops longi-caudatus, Murch.; Dalmanites longicaudatus, enmienda de Font y Sagué. Notas Inst. geol. Esp. No. 13 1944: 5-6 2 pls.

Samsonowicz, J. (1). On the Holmia-Fauna in the Cambrian of the Anticlinorium of Klimontów. Bull. Acad. polon. Sci. [sci. chim., géol., géogr.] 7 6 1959: 447–452 pls. i–iii text-fig. 1.

Samsonowicz, J. (2). On Strenuaeva from Lower Cambrian in Klimontów Anticlinorium. Bull. Acad. polon. Sci. [sci. chim., géol., géogr.] 7 7 1959: 521-524 pl. i.

Samsonowicz, J. (3). On Strenuella and Germaropyge from the Lower Cambrian in the Klimontów Anticlinorium. Bull. Acad. polon. Sci. [sci. chim., géol., géogr.] 7 7 1959; 525-529 pls. i-ii.

Schmidt, H. Gefälschte Versteinerungen. Natur u. Volk 89 1 1959: 27-29 text-figs. 1-4.

Schmidt, H. see Harrington, H. J.

**Sdzuy,** K. Die unterkambrische Trilobiten-Familie Dolerolenidae. Senckenbergiana leth. **40** 5–6 1959 : 389–407 pl. i text-fig. 1.

Sdzuy, K. see Harrington, H. J.

Seilacher, A. Vom Leben der Trilobiten. Naturwissenschaften 46 12 1959: 389-393 text-figs. 1-5.

Semikhatov, M. A. [Lower Cambrian of the Kansk-Angara depression.] Bull. Soc. nat. Moscou (ns) 64 [Geol. 24 2] 1959: 91-109 text-figs. 1-3 [In Russian, with English summary p. 109].

Shackleton, R. M. The Stratigraphy of the Moel Hebog District between Snowdon and Tremadoc. Liv. Manchr geol. J. 2 2 1959: 216-252 pl. xvi (map) text-figs. 1-3 table [Trilobita det. W. F. Whittard, with A note on the Caradoc shelly faunas by J. C. Harper pp. 249-250].

Shaw, A. B. Quantitative trilobite studies III. Proliostracus strenuelliformis Poulsen, 1932. J. Paleont. 33 3 1959: 474-487 text-figs. 1-15.

Shcheglova, O. S. see Miroshnikov, L. D.

Shelkovnikov, A. D. see Anatol'eva, A. I.

Sinclair, G. W. see Thorsteinsson, R.

Sivoy, A. G. see Anatol'eva, A. I.

Snaidr, M. see Prantl, F.

**Strad,** V. (1). První nález trilobita v grygovských vápencich. Zprávy SLUKO, Olomouc No. 48 1955 : 7–8 text-fig.

Strnad, V. (2). Harpes neogracilis R. et E. Richter, 1924—nový trilobit v československém devonu. Zpvávy KVM [Zprávy SLUKO], Olomoue No. 64 1956: 104–105 pl.

Strnad, V. (3). Výskyt trilobita Chasmops odini (Eichwald, 1840) z baltského ordoviku v souvoích v Olomouckém kraji. Zprávy KVM [Zprávy SLUKO], Olomouc No. 66 1956: 130-131 text-fig.

Strnad, V. (4). Dechenella (Basidechenella) dombroviensis (Gürich, 1896)—další nový tribolit z moravského devonu. Zprávy KVM [Zprávy SLUKO], Olomouc No. 66 1956: 136.

Strnad, V. (5). Největší rozšíření plistocenního ledovce a další nový trilobit pro ČSR ze sandrových písků. Zprávy KVM [Zprávy SLUKO], Olomouc No. 70 1957: 39-40 text-fig.

Strnad, V. (6). O Chabičově, důležitém moravském nalezišti devonských zkamenělin. Zprávy KVM [Zprávy SLUKO], Olomouc No. 72 1957: 71–72 2 text-figs.

Strnad, V. (7). Nomenklatorická pravidla a severský trilobit od Vidnavy. Zprávy KVM [Zprávy SLUKO], Olomoue No. 73 1957: 91.

Strnad, V. (8). Trilobiti z glaciofluviálních štěrkopísků od Vidnavy. Čas. Min. Geol., Prague 3 3 1958: 318–326 pls. xxv-xxvii [German summary pp. 326–327].

Struve, W. (1). Beiträge zur Kenntnis der Phacopacea (Trilobita), 4: Volkops volki n.g., n.sp., ein Phacopinae aus dem deutschen Ordovizium. Senekenbergiana leth. 40 1–2 1959: 29–42 pls. i-ii.

Struve, W. (2). Beiträge zur Kenntnis der Phacopacea (Trilobita), 5: Gourdonia destombesi n. sp. (Asteropyginae) aus dem Mittel-Devon von S-Frankreich. Senckenbergiana leth. 40 1-2 1959: 47-50 text-figs. 1-3.

Struve, W. see Harrington, H. J.

Stubblefield, C. J. (1). [Report of the] Palaeontological Department. Summ. Progr. geol. Surv. G. B. (for 1951) 1953: 56-59.

Stubblefield, C. J. (2). Lexique stratigraphique international I Europe fasc. 3s Angleterre, Pays de Galles, Écosse III Cambrien. Centre National de la Recherche Scientifique, Paris 1958 [1959]: 95 pp.

Stubblefield, C. J. (3). Evolution in trilobites. Quart. J. geol. Soc. Lond. 115 2 1959: 145-162 text-figs. 1-2.

Stubblefield, C. J. see Harrington, H. J.

[1

Clarch

Tre

silte

calc

axi

Ins

Fe

str Tr

En

A A

M

M

Suñer Coma, E. & Fernández de Villalta, J. Un nuevo trilobites en Camprodón (Gerona). Mem. y Com. Inst. geol. Barcelona 16 1957: 19-31 text-figs. 1-5.

Suvorova, N. P. (1). [On the morphology and systematic position of the trilobites of the family Granularidae.] Paleont. Zhurnal, Moscow 1959 2: 101-106 text-figs. 1-3 [In Russian].

Suvorova, N. P. (2). [New trilobites of the superfamilies Corynexochoidea and Redlichioidea of the Lena stage of the Lower Cambrian of Yakutia.] Paleont. Zhurnal, Moscow 1959 3: 66-77 pl. vi textfigs. 1-5 [In Russian].

Theokritoff, G. Taconic sequence in northern Washington County, New York [Abstract]. Bull. geol. Soc. Amer. 70 12 1959: 1686–1687.

Thorsteinsson, R. Cornwallis and Little Cornwallis Islands, District of Franklin, Northwest Territories. Mem. geol. Surv. Canada No. 294 1958: v+134 pp. 8 pls. 7 tables 5 figs. map [Trilobita det. A. E. Wilson, G. W. Sinelair, D. J. McLaren].

Tjernvik, T. E. The Tremadocian Beds at Flagabro in South-Eastern Scania (Sweden). Geol. Fören. Stockh. Förh. 80 3 1958: 259–276 pl. iii text-figs. 1-6

Tripp, R. [P.] see Harrington, H. J.

Vanék, J. (1). Nové zjištěné rody trilobitů v Barrandienu. Čas. nár. Mus. [Přír.] 127 2 1958 : 215.

Vanék, J. (2). Čeled' Lichaidae Hawle et Corda, 1847 ze středočeského staršího paleozoika (Trilobitae). Bohemia Centralis, Prague [A] 1 3 1959: 79–168 pls. i-xii text-figs. A-F, 1–43 [English summary pp. 156– 1591

Vanék, J. (3). Fauna příležitostných odkryvů na území Velké Prahy. Zprávy geol. výzkum, for 1959:

Veber, V. N. see Lesnikova, A. F.

Walmsley, V. G. The geology of the Usk inlier (Monmouthshire). Quart. J. geol. Soc. Lond. 114 4 (for 1958) 1959: 483-516 pl. xxii text-figs. 1-8 table I.

Walmsley, V. G. see Holland, C. H.

Weber, V. N. see Lesnikova, A. F.

Weir, J. A. Ashgillian Trilobites from Co. Clare, Ireland. Palaeontology 1 4 1959: 369–383 pls. lxii-lxiii.

Weller, J. M. see Harrington, H. J.

Whittard, W. F. see Shackleton, R. M.

Whittington, H. B. Silicified Middle Ordovician Trilobites: Remopleurididae, Trinucleidae, Raphiophoridae, Endymioniidae. Bull. Mus. comp. Zool. Harv. 121 8 1959: 371-496 pls. i-xxxvi text-figs. 1-8.

Whittington, H. B. see Harrington, H. J.

Whittington, H. B. see Kindle, C. H.

Williams, M. Y. The Geological History of Churchill, Manitoba. Western Miner, Vancouver 21 6 1948: 39-42 text-figs.

Wilson, A. E. see Thorsteinsson, R.

Yanagisawa, I. [On a Discovery of Trilobite (Permian) from the Takakura Yama Group, Abukuma Mountainland.] J. geol. Soc. Japan 64 No. 751 1958: 207 text-fig. [In Japanese].

Yanishevsky, M. E. see Lesnikova, A. F.

Zhuravleva, I. T. & Repina, L. N. [Generic complexes of trilobites and archaeocyathids of the Lower Cambrian of the Altai-Sayan region.] C. R. Acad. Sci. U.R.S.S. (ns) 129 1 1959: 181–183 pls. i–ii (charts) [In Russian].

Zhuravleva, I. T., Repina, L. N. & Khomentovsky, V. V. (1). [Biostratigraphy of the Lower Cambrian of the folded frame of the Minusinsk depression.] Bull. Soc. nat. Moscou (ns) 64 [Geol. 34 2] 1959: 67-90 text-figs. 1-2 [In Russian with English summary p. 90].

Zhuravleva, I. T., Repina, L. N. & Khomentovsky, V. V. (2). [Lower Cambrian horizons of the Shory highlands.] C. R. Acad. Sci. U.R.S.S. (ns) 128 5 1959: 1030-1033 text-figs. 1-2 [In Russian].

Zhuravleva, Z. A. (1). [On the stratigraphy of the Cambrian of the middle and lower reaches of the river Olekma.] C. R. Acad. Sci. U.R.S.S. (ns) 128 4 1959: 798-799 [Trilobita det. L. N. Repina] [In Russian].

Zhuravleva, Z. A. (2). [On the stratigraphy of the ancient sedimentary deposits of the basin of the middle reaches of the river Chara.] C. R. Acad. Sci. U.R.S.S. (ns) 129 6 1959: 1370-1373 [Trilobita det. N. V. Pokrovskaya] [In Russian].

#### II.—SUBJECT INDEX

#### GENERAL LITERATURE AND HISTORY

General works.—Treatise, Harrington et al.;
Trilobite evolution, Stubblefield (3); Dolerolenidae,
Sdzuy; Lichidae of Czechoslovakia, Vaněk (2);
Little atlas of fossils, Bodylevsky; Correlation tables
for Precambrian and Palaeozoic of Kazakhstan,
Borukaev et al.; Cambrian and Ordovician limestones
of Sweden, Hadding; Stratigraphy of W. Australia,
McWhae et al.

Popular works.—Palaeozoic of Estonia, Kaljo (1); Fossil collecting, Kaljo (2).

Reference books.—Stratigraphic dictionary—Cambrian of Britain, Stubblefield (2).

Obituary.—R. & E. Richter, Gómez de Llarena, Kegel.

Abstracts.—Fritz, Jones, Kauffman, Robison, Theokritoff.

Reviews.—Last decade of palaeontology, Erben; Palaeontology and stratigraphy in Sweden, Regnéll; Šnajdr 1958, Prantl. Check lists.—Lower Cambrian trilobites and archaeocyathids of Altai-Sayan, Zhuravleva & Repina; Tremadoc trilobites of Norway, Henningsmoen.

Preservation.—In Elrathia kingii preserved in siltstone the test is replaced by cryptocrystalline calcite, cone-in-cone incrustations occur on ventral surfaces, and there may be cracks parallel to thoracic axial furrows which simulate nodes on the axis, Bright.

Museum Collections,—Cambrian fossils of the Instituto Geológico y Minero de España, Badillo.

Forgeries.-Forged Phacops latifrons, Schmidt.

Revision of fauna.—Revision of faunal lists of Ferronnière 1920-22, Pillet (2).

Revision of nomenclature.—Validation of Entomostracites punctatus Wahlenberg 1821 (by suppression of Trilobus p. Brünnich 1781) and designation as type of Encrinurus Emmrich 1844; designation of Asaphus hausmanni Brongniart 1822 as type of Odontochile Hawle & Corda 1847, Opinion 537; Designation of Asaphus angustifrons Dalman as type of Ptychopyge Angelin, Opinion 538; Kendallia Berg 1953 non Evermann & Shaw 1927 renamed Kendallina, Mitchellia Vogdes 1917 non de Koninek 1877 renamed Mitchellaspis, Harrington et al.

Additions to Official Lists,—Encrinuridae Angelin 1854, Encrinurus Emmrich 1844, E. punctatus (Wahlenberg 1821), Odontochile Hawle & Corda 1847, O. hausmanni (Brongniart 1822), Opinion 537; Ptychopyge Angelin 1854, P. angustifrons (Dalman 1827), Opinion 538.

Designation of type species.—Encrinurus Emmrich 1844—designation of Entomostracites punctatus Wahlenberg 1821, Odontochile Hawle & Corda 1847—Asaphus hausmanni Brongniart 1822, Opinion 537; Ptychopyge Angelin 1854—designation of Asaphus angustifrons Dalman 1827, Opinion 538.

Selection of neotypes,—Phacopidella (Dienstina) liopyga Reinh. Richter, Pfeiffer; Trochurus speciosus Beyrich 1845, Vaněk (2).

Selection of lectotypes.—Acanthopyge haueri (Barrande 1846), A. leuchtenbergii, A. speciosa, A. pulchra, Corydocephalus flabellatus, C. interjectus, C. propinquus, C. verrucosus, Dicranogmus pustulatus, Dicranopeltis aspera, D. granulosa Hawle & Corda 1847 spp., Hemiarges ambiguus (B. 1846), Lichas incola B. 1856, Platylichas kloučeki (Růžička 1926), Uralichas avus (B. 1872), U. giganteus (Klouček 1916), Vaněk (2); Alokistocare althea Walcott, Anoria tontoensis (W.), Glossopleura mekee Resser; Ampyx camurus Raymond 1925, Whittington.

#### STRUCTURE

Terminology.—Term genicranium for dorsal cephalic shield of hypoparian forms and Olenellids, separated by marginal (or just supramarginal in Conocoryphids) suture from ventral doublural plate; rostri-hypostomal plate for fused rostrum and hypostome, librigenal plate for conjoined free cheeks, Henningsmoen; Several morphological terms introduced for Remopleuridids, Whittington.

Glabellar furrows.—Furrows in Remopleurides are not depressions in the external surface but lines along which the test is thinner, Whittington.

Cephalic spines.—Pairs of marginal spines in protaspides of several genera; frontal glabellar spine of Ampyx appears suddenly in meraspis O, in Ampyxina it is gradually reduced in later development, Whittington.

Genal spines.—Present in Ordovician Volkops (Phacopinae), Struve (1).

Genal caeca.—Cheek wrinkles on cephala of Agnostids (illustrated in *Ptychagnostus*) so described and interpreted as hepatic glands, **Öpik** (1).

Alae of Raphiophoridae considered not parts of the glabella, Whittington.

Eyes.—Present in Raphiophorid Ampyxinella, Koroleva.

Ventral sutures.—Median suture separating free cheeks in *Leiocoryphe*, Rasetti; No median or connective sutures in Raphiophoridae, Whittington.

Hypostome.—Comparison of hypostomes of Remopleurides and Ceraurus suggests hypothesis that hypostome of R. corresponds only to posterior part of that of C., the anterior part of C. (including wing process) being represented by part of cephalic doublure in R., the hypostomal sutures in the generabeing not homologous; hypostomes of Cryptolithus tesselatus and of several Raphiophorids described and suggested not to have been in sutural contact with cephalic doublure but held closely up under glabella, Whittington.

Numbers of thoracic segments.—A small percentage (about 2-5) of "adult" Elrathia kingii specimens show up to three fewer than the normal thirteen thoracic segments, Bright.

Pygidium.—'Open' pygidium (i.e. without posterior axial doublure) described in Saukianda and considered in this case not a sexually dimorphic character, Sazuy.

Ornamentation.—5% of Elrathia kingii specimens show granulation, Bright.

Pits.—Flat bases of pits of upper and lower lamellae of fringe of Cryptolithus tesselatus are in contact, opposing pits being apparently connected by a minute hole in each base, Whittington.

Integuments.—Several sections of limestones containing trilobite tests, Hadding.

Alimentary system.—Genal caeca interpreted as hepatic glands; inferred relations of glands and stomach in Agnostids; intestinal appendages stated to occur on thoracic segments and pygidia of Ptychopariids, Öpik (1); Radiating punctae on internal moulds of cheeks and pygidial pleural regions of Pseudagnostus prolongus interpreted as caeca, Lochman & Hu.

Musculature.—Glabellar and pygidial muscle scars in Pseudagnostus prolongus, Lochman & Hu.

#### PHYSIOLOGY AND SEX

Enrolment.—Suggested that Remopleurides simulus could not completely enrol, Whittington.

Sexual dimorphism.—Three suggested cases in Upper Cambrian forms in which the inferred female has an occipital spine and different glabellar convexity—Arapahoia neihartensis & A. aspinosa, Cedarina cordillerae & C. prima, Minkella americana & Q. & J. Lochman & Hu: ; Dimorphism (narrow/brosd) in mature pygidia of Weberides mucronatus analysed biometrically and considered sexual, Rehof & Rehofová; Remopleurides eximius and R. simulus may be dimorphs, Whittington; 'Open' pygidium considered not a sexually dimorphic character in Saukianda, Sázuv.

#### DEVELOPMENT

Ontogeny.—Protaspides and later developmental aggs of spp. of Remopleurides (protaspis enrolled, with fixigenal and anterior cephalic spines), Cryptolithus, Tretaspis (referred protaspis with 3 pairs of marginal spines), Raphiophoridae (referred protaspis of Ampyx has 2 pairs of marginal spines; frontal glabellar spine appearing in meraspis O), Whittington; Immature stages of Idahoia, Minkella and Ptychaspis, Lochman & Hu; Immature stages of Prosaukia corrugata (smallest observed cranidium is only 0.3 mm. long), Rasetti.

Ecdysis.—Sample of 339 Elrathia kingii does not show instars, Bright.

#### **EVOLUTION AND CLASSIFICATION**

Evolution.—General discussion of trilobite evolution; suggested examples of neoteny and gerontism, Stubblefield (3).

Phylogeny.—Possible relations of Granulariidae and Asaphoidea, Suvorova (1).

Abundance.—Charts of subfamily abundance, Stubblefield (3).

Homeomorphy.—Several examples discussed, Stubblefield (3),

Biometry.—All possible combinations in pairs of 6 dimensions on 4 Poulsen 1932 spp. of Proliostracus show rectilinear relations; concluded that spp. represent growth-stages of a single sp. P. strenuelliformis of which the others are considered synonyms, Shaw; Study of sample of 339 Elrathia kingii, rectilinear relations between dimensions plotted, Bright; Pygidial dimorphism (narrow/broad) in Weberides mucronatus analysed, Rehoř & Rehořová.

#### ECOLOGY AND HABITS

**Ecology.**—Ecology of a *Ptychaspis* faunule discussed: indigenous unsorted benthos, more or less extraneous mechanically sorted benthos, and plankton distinguished, **Lochman & Hu**.

Habits.—General account of crawling, grazing, burrowing and tunnelling; spinous forms (as Acidaspis) considered too large to be planktonic, Seilacher; Elrathia kingii inferred to have inhabited calm shallow water on a calcareous silty bottom, Bright; Kootenia and Solemopleurella restricted to limestones, Elrathia to fine sands and silts, Alokistocare and Glossopleura more tolerant, McKee.

Trails.—Walking trail possibly attributable to trilobites, Opik (2); McKee.

Migration.—Lower Cambrian faunal affinities between Poland, Britain and Western Mediterranean, Samsonowicz (3).

#### DISTRIBUTION AND FAUNAS

Cambrian.—Britain, Stubblefield (2); Canada (Rooky Mts.), North & Henderson; Spain, Badillo; U.S.S.R. (N. W. Siberia), Miroshnikov et al.; U.S.S.R. (W. Siberia), Petrakov.

Cambrian (Lower).—Greenland (East), Shaw; Poland (Holy Cross Mts.), Samsonowicz (1), (2), (3); Spain (León, Andalusia), Sdzuy; U.S.A. (New York), Theokritoff; U.S.S.R. (Altai), Zhuravleva et al. (2); U.S.S.R. (Siberia), Z.A. Zhuravleva (1), (2); U.S.S.R. (S. Siberia), Semikhatov; U.S.S.R. (Tuva, Sayan etc., Yakutia), Pokrovskaya; U.S.S.R. (Yakutia), Suvorova (2).

Cambrian (Lower & Middle).—Australia (S. Australia), Glaessner & Parkin; U.S.A. (Arizona), McKee, Resser; U.S.S.R. (Altai-Sayan), Zhuravleva & Repina; U.S.S.R. (Sayan), Anatol'eva & Shelkovnikov; U.S.S.R. (S. Siberia), Zhuravleva et al. (1).

Cambrian (Middle). — Argentina (Mendoza), Rusconi; Poland (Holy Cross Mts.), Orlowski (1), (2), (3); U.S.A. (Montana), Kaufiman; U.S.A. (Utah), Bright; U.S.S.R. (N. Siberia), Lazarenko.

Cambrian (Middle & Upper).—Canada (W. Newfoundland), Kindle & Whittington; China (North), Manchuria, Chang.

Cambrian (Upper).—Britain (N. Wales), Shackleton, Stubblefield (1); U.S.A. (Colorado), Berg & Ross; U.S.A. (Idaho, Texas), Lochman & Hu; U.S.A. (Md, Pa, Va), Canada (Quebee), Rasetti; U.S.A. (Utah), Robison; U.S.S.R. (Kazakhstan), Ivshin.

Ordovician.—China, Lu; Czechoslovakia (boulders), Strnad (3), (5), (7), (8); Germany (Thuringia), Struve (1); Nepal, Bordet et al.; U.S.S.R., Lesnikova & Weber; U.S.S.R. (N. W. Siberia), Miroshnikov et al.

Ordovician (Lower).—Canada (Rocky Mts.), North & Henderson; Norway, Henningsmoen; Sweden (Scania), Tjernvik; U.S.A. (Colorado), Berg & Ross; U.S.S.R. (Baltic, Kazakhstan), Balashova & Balashov.

Ordovician (Lower & Middle).—Britain (N. Wales), Shackleton; U.S.S.R. (W. Siberia), Petrakov.

Ordovician (Middle).—Canada (Ontario), Colquboun; U.S.A. (mainly Virginia), Britain (Scotland), Whittington.

Ordovician (Middle & Upper).—Britain (N. England), Dean (1); Canada (Arctic), Thorsteinsson; Czechoslovakia, Yaněk (3); U.S.S.R. (Arctic), Balashova (1), (2); U.S.S.R. (Kazakhstan), Koroleva.

Ordovician (Upper).—Britain (N. England, Wales), Dean (2); Czechoslovakia, Vaněk (1); Eire, Weir; Estonia, Männil.

Ordovician or Silurian.—Canada (Manitoba), Williams.

Silurian,—Canada (Arctic), Thorsteinsson; Czechoslovakia (boulders), Strnad (8); U.S.S.R., Lesnikova & Weber; U.S.S.R. (N. W. Siberia), Miroshnikov et al. Silu Suñer (Arcti Silu

[195

Silu Walm Silu et al. Vaněl

Der Franc Suñer

> Maks De Paul

Cana (1); Fran (Nev

> Pola Ca Ca

Pole Reh P

† i fos

ni

p.

of fi

Silurian (Lower).—Spain (Gerona), Sampelayo; Suñer Coma & Fernández de Villalta; U.S.S.R. (Arctic), Balashova (1), (2).

Silurian (Middle).-Japan (Kyushu), Hamada.

Silurian (Middle & Upper).—Britain (Wales),

Silurian (Upper).—Britain (W. England), Holland et al.; Britain (Wales), Holland; Czechoslovakia, Vaněk (1), (2).

Devonian.—Canada (Arctic), Thorsteinsson.

Devonian (Lower).—Czechoslovakia, Vaněk (2); France (Anjou), Pillet (1), (2); Spain (Gerona), Suñer Coma & Fernández de Villalta.

Devonian (? Lower).—U.S.S.R. (Sikhote Alin), Maksimova & Organova,

Devonian (Lower & Middle).—Germany (Eifel), Paulus.

Devonian (Middle).—Belgium (Liège), Monseur; Canada (Arctic), McLaren; Czechoslovakia, Strnad (1); Czechoslovakia (Silesia), Strnad (2), (4), (6); France (Haute Garonne), Struve (2); U.S.A. (Nevada), Jones.

Devonian (Upper).—Britain (S. W. England), Dearman & Butcher; Germany (Thuringia), Pfeiffer; Poland (Silesia), Lewowicki.

Carboniferous.-U.S.S.R. (European), Ivanova.

Carboniferous (Lower).—Britain (Eire), Caldwell; Poland (Silesia), Lewowicki.

Carboniferous (Upper).—Czechoslovakia (Silesia), Rehoř & Rehořová.

Permian.-Japan (Fukushima), Yanagisawa.

#### III.—SYSTEMATIC INDEX

N.B.—Since all the species of the group are fossil, the † used elsewhere in the Zoological Record to indicate fossils is here omitted.

Abadiella Hupé 1952 [sic] discussed and referred to Doleroleninae p. 398, **Sdzuy.** 

Abadiellidae Hupé 1952 [sic] a synonym of Doleroleninae p. 398, **Sdzuy**.

Acanthaloma sp. p. 78, Lu.

Acanthaloma sp. p. 103, Pillet (2).

Acanthoparypha chiropyga p. 384, A. perforata p. 385 ff., A. sp. p. 387, Whittington.

Acanthopyge Hawle et Corda 1847 discussed p. 119, A. haueri (Barrande 1846) lectotype selected and figured excluding records of Maurer 1880 and Novák 1890 and with synonym A. leuchtenbergii H. & C. of which lectotype selected and figured p. 120 pl. v figs. 1-7, pl. vi figs. 3-6, text-figs. 14-17 [A. aff. h. of Chlupåć & Vaněk 1957 referred to Lobopyge docekali sp. n. p. 125], A. parva parva (Barrande 1846) holotype [= Lichas h. of B. 1852 partim (pl. 28 fig. 38)] refigured with synonym L. parvula Novák 1890

(excluding records of parvula by Richter 1909 and cf. p. of R. & E. R. 1917) p. 125 pl. vi figs. 1-2; A. pulchra pl. ix fig. 9 and A. speciosa pl. viii fig. 10 H. & C. spp. lectotypes chosen and figured and synonyms of Trochurus speciosus p. 147, Vanék (2).

Acaste downingiae (Murchison) p. 514, Walmsley.

Acaste (Acastella) rouaulti (de T. et L.) p. 18 ff., Pillet (1).

Acastella see Acaste.

Acastinae see Duftonia gen. n.

Acastoides cf. henni posthuma (R. and E. Richter) p. 342, Paulus.

" Acidaspis" cf. asteroidea Reed p. 218, Dean (1).

"Acidaspis" ulrichi Bassler so referred and stated to be Upper Cambrian in age and to have conjoined free cheeks p. 393 pl. li fig. 27, Rasetti.

" Acidaspis" ulrichi p. 17, Kindle & Whittington.

Acidaspis sp. p. 514, Walmsley.

Acrocephalaspis gen. n. [no family assigned] p. 107 for type A. fidus sp. n. p. 109 pl. i figs. 1, 2, 2a Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Acrocephalites sp. n. [undescr.] p. 360, Miroshnikov et al.

Acrocephalops see Bolaspididae fam. n.

Acrocephalops & arizonaensis sp. n. p. 203 pl. xxvi fig. 11 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg, Instn No. 563 1945.

Acrolichas punctatus Weber 1948 p. 304 pl. lxxx figs. 13a-14, Lesnikova & Weber.

Actinopeltis caudata n. sp. [nom. nud.] p. 74 Ordovician China, Lu.

Adiaksis [nom. nud.] sp. p. 1032, Zhuravleva et al. (2).

Aeglina rediviva record of Baily 1862 referred to Phylacops bituberculatus sp. n. p. 377, Weir.

Agnostacea discussed p. 152, Stubblefield (3).

Agnostids—genal caeca described on cephala and similar structures on pygidia p. 1750, Öpik (1).

Agnostids p. 1686, Theokritoff.

Agnostus bidens Meek p. 84, Bright.

Agnostus calvus (Lake) p. 223, A. sp. p. 222, Shackleton.

Agnostus pisiformis Linn. p. 117 (pl. 43) fig., A. integer Beyr. p. 118 (pl. 44) fig., Badillo.

" Agnostus" sp. p. 47, Vaněk (3).

Agnostus sp. undet. p. 380 pl. li fig. 10, Rasetti.

Agraulus [sic] (Arionellus) ceticephalus Barr. p. 109 (pl. 35) fig., Badillo.

Alacealus Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

[19

Ap

viciai

Whit

Caml

An

A

A

A

A

Pty

A. J

A

Lu.

Th

an [q

[ty

pe W

Bals

Pale

1944

in L

Alataja Poletaeva [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Albertella schenki sp. n. p. 195 pl. xx fig. 18 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Aldonaia Lerm. 1940 discussed p. 89, A. cf. ornata L. p. 90 pl. viii fig. 16, A. shanganica sp. n. p. 91 pl. viii figs. 14, 17, A. punctuosa sp. n. p. 93 pl. viii figs. 3, 6 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud, geol. Inst. S.S.R. 27 1959.

Aldonaidae see Eleganolimba and Tuvanella genn. n.

Alokistocare althea Walcott lectotype selected and figured p. 203 pl. xxii figs. 6-11, A. sp. undet. p. 203 pl. xxii fig. 5, Resser,

Alokistocare elongatum Poulsen discussed p. 9,

Amecephalina range in Altai-Sayan pl. i, Zhuravleva & Repina.

Amecephalina sp. p. 1033, Zhuravleva et al. (2).

Amecephalus mendozanus (Rusc.) discussed p. 9, Rusconi.

Amphilichas batchaticus Weber 1928 p. 304 pl. lxxxii figs. 3a-c, Lesnikova & Weber.

Amphilichas holmi (Schm.) p. 7, A. lineatus Ang. p. 11, Mannil,

Amphilichas pandus p. 384, Whittington.

Ampyx depressus (Angelin) var. otradnica var. nov. pl. i figs. 20-23, A. nordicus sp. nov. pl. vi figs. 1-5, A. borealicus sp. nov. pl. vi figs. 8-9, Balashova (1).

Ampyx nasutus Dalman 1826 p. 276 pl. lxv figs. 9, 10a-b. Lesnikova & Weber.

Ampyx rugosus Kolova type of Ampyxinella gen. n. p. 1314, Koroleva.

Ampyx virginiensis Cooper 1953 morphology and ontogeny (including protaspis) described p. 465 pl. xxix figs. 1-16, pl. xxx figs. 1-14, 16-17, 20-30, pl. xxxi figs. 1-2 [? attribution p. 464], 3-32, text-figs. 7A-B (reconstrs.), A. camurus Raymond 1925 lectotype selected [pl. expl.] and figured p. 469 pl. xxx figs. 15, 18-19. Whittington.

Ampyxina Ulrich 1922 discussed, A. bellatula (Savage 1917) holotype refigured p. 481 pl. xxxiv figs. 1-2, A. powelli (Raymond 1920) morphology and ontogeny p. 482 pl. xxxiv figs. 3-13, pl. xxxv figs. 1-25, A. lanceola 5p. n. including A. elegans Cooper 1953 partim (pl. 4 figs. 10-11) Ordovician (Middle) U.S.A. (Virginia) p. 486 pl. xxxiv figs. 14-28, pl. xxxv figs. 26-35, A. aldonensis (Reed 1935) p. 487 pl. xxvii figs. 10-11, 13; A. elegans C. 1953 restr. [non pl. 4 figs. 10-11 to A. l., figs. 13, 17 to Tretaspis sagenosus] referred to Raymondella p. 488, as is A. sp. C. 1953 p. 490, Whittington Bull. Mus. comp. Zool. Harv. 121 8 1959.

Ampyxina biloba Chugaeva referred to Ampyxinella gen. n. p. 1315, Koroleva.

Ampyxina taimyrica sp. nov. pl. vi figs. 10-14, Balashova (1).

Ampyxinella gen. n. [Raphiophoridae (Ampyxinellinae)] p. 1314 for type Ampyx rugosus Kolova 1936 pl. figs. la-6, text-fig. 4a, A. balaschova sp. n. [nom. nud.] pl. figs. ls, ε, A. ruslani sp. n. [nom. nud.] pl. fig. lo, A. rugosa var. nanusi var. nov. [nom. nud.] pl. fig. le, A. biloba (Chugaeva) for Ampyxina b. C. p. 1315 Ordovician (Middle & Upper) U.S.S.R. (Kazakhstan), Koroleva C. R. Acad. Sci. U.R.S.S. (ns) 124 6 1959.

Ampyxinellinae [misspelt Alpyxiinellinae at descr.] subfam. n. [Raphiophoridae] for Ampyxinella gen. n. p. 1314, Koroleva C. R. Acad. Sci. U.R.S.S. (ns) 124 6 1959.

Anabaraspis sp. p. 5, Lazarenko.

 $An a doxides \ {\bf Matthew} \ 1899 \ {\bf a} \ {\bf synonym} \ {\bf of} \ {\it Metadoxides} \\ {\bf p.} \ 397, \ {\bf Sdzuy.}$ 

Andrarinidae see Olenina.

Anomocare Angelin 1852—attention drawn to Vogdes' designation of A. aculeatum (Angelin) as type p. 287, STUBBLEFIELD in Harrington et al.

Anomocare see Szeaspis gen. n.

Anomocare parvula Weller reference to Bowmania unconvincing p. 396, Rasetti.

Anomocarella tumida Endo 1944 apparently referred to Metanomocarella gen. n. q. v. [though not as type] and Aojia tumida [Resser &] Endo 1937 may be a synonym p. 218, Chang.

Anomocarellidae see Luia gen. n.

Anomocaridae see Olenina.

Anomocarina sp. indet. p. 359, Miroshnikov et al.

Anoria tontoensis (Walcott) lectotype selected and figured p. 197 pl. xxii figs. 7-15, pl. xxii figs. 1-4, Resser.

Antagmellus [Antagnellus on pl.] Lermontova [nom. nud.] range in Altai-Sayan p. 183 pl. i, Zhuravleva & Repina.

Antagmidae see Lermontoviella gen. n.

Antagmus arizonaensis sp. n. p. 205 pl. xix figs. 27–28 Cambrian (Lower) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Aojia tumida [Resser &] Endo 1937 may belong to Anomocarella tumida E. 1944 which is referred to Metanomocarella gen. n. [quod vide] p. 218, A. longispina R. & E. attributed pygidium (1937 pl. 47 fig. 2) belongs to Taitzuia p. 223, Chang.

Apatocephalus assai Weber 1932 p. 275 pl. lxv fig. 2, Lesnikova & Weber.

Apatokephaloides minor sp. n. p. 387 pl. lv figs. 29-32, A. macrops sp. n. p. 388 pl. lv figs. 24-28 Cambrian (Upper) U.S.A. (Maryland), A. sp. undet. p. 378, Rasetti J. Paleont. 33 3 1959.

Apatokephalus serratus (Boeck) listed p. 174, Henningsmoen.

Apatokephalus serratus (Boeck) p. 274, Tjernvik.

Apatokephalus serratus Sars et Boeck p. 144, Balashova & Balashov.

al.

36

m.

1.1

1.]

b.

R.

r.1

24

es

to

ia

e]

d

i,

0

Apatokephalus yini n. sp. [nom. nud.] p. 88 Ordovician China, Lu.

Apianurus barbatus, A. glaber, A. sp. ind. p. 385 ff., Whittington.

Aposolenopleura plicata sp. n. p. 395 pl. li figs. 19–22 Cambrian (Upper) U.S.A. (Maryland), Rasetti J. Paleont. 33 3 1959.

Arapahoia aspinosa & A. neihartensis Lochman 1944 spp. considered ♂ & ♀ of one sp. p. 415, Lochman in Lochman & Hu.

Arionellus see Agraulus.

Aristoharpes taimyricus sp. nov. pl. v figs. 20–21, Balashova (1).

Aristoharpes sp. p. 1548, Bordet et al.

Asaphellus alimbeticus Bal. [not seen] p. 144; Ptychopyge? inostranzewi Lamansky 1905 belongs to A. p. 145, Balashova & Balashov.

Asaphellus homfrayi (Salter) p. 223, Shackleton.

Asaphellus inflatus n. sp. p. 33, A. nanchuangensis n. sp. p. 37 [nom. nuda] Ordovician (Lower) China, Lu.

Asaphid p. 233 ff., Shackleton.

Asaphidae see Niobinae, Promegalaspidinae & Thysanopyginae  $\operatorname{subfamm}$ .

Asaphids p. 384 ff., Whittington.

Asaphiscus—spp. of Resser & Endo 1937 discussed and the following referred to Liaoyangaspis gen. n. [quod vide]: bassleri (with synonym kobayashii) [type of L.], valcotti restr. (excluding R. & E. pl. 43 fig. 6 to L. endo: sp. n., & Lu 1957 to L. vanyi sp. n.), peiensis, & tsutsumii (E. 1937) p. 220; A. iddingsi Walcott referred to Szeaspis gen. n., A. cf. i. of Chang 1957 to S. reticulatus sp. n. p. 230, Chang.

Asaphiscus see Szeaspis gen. n.

Asaphiscus wheeleri Meek p. 84, Bright.

Asaphoidea—relations to Granulariidae & Dikellocephaloidea p. 106, Suvorova (1).

Asaphus Brongniart specific key p. 282, A. platyurus Angelin 1854 p. 282 pl. lxxii figs. 4–5, A. delphinus Lawrow 1856 p. 282 pl. lxxii figs. 2–3, A. kovalewskii L. 1856 p. 283 pl. lxx fig. 1, lxxii fig. 4, A. cornutus Pander 1830 p. 283 pl. lxx fig. 3, A. cornutus var. punctatus Lesnikova subsp. n. Ordovician (Middle) U.S.S.R. (Leningrad) p. 283 pl. lxx figs. 2a–b, A. c. var. holmi Schmidt 1901 p. 283, A. devexus Eichwald 1840 p. 284 pl. lxiv figs. 1–5, A. bröggeri S. 1898 p. 284 pl. lxvii figs. 1–2, A. priscus Lamansky 1905 [diagnosis & illustration] p. 284 pl. lxix figs. 1a–b, A. expansus Dalman 1827 p. 285 pl. lxviii figs. 3–4, A. lepidurus Nieszkowski 1859 p. 285 pl. lxviii figs. 3–4, A. lamanskii S. 1901 p. 285 pl. lxviii figs. 8–9, pl. lxix figs. 2, A. raniceps D. 1827 p. 285 pl. lxviii figs. 5, pl. lxvii figs. 3a–b, A. pachyophtalmus S. 1898 p. 286 pl. lxviii figs. 10–11, A. ornatus Pompecki 1890 p. 286 pl. lxvii figs. 1a–c, A. eichwaldi S. 1901 p. 286 pl. lxviii figs. 12–13, A. latus Pander 1830 p. 287 pl. lxxi figs. 1a–b, A. la evissimus S. 1898 p. 287 pl. lxxii figs. 1a–b, A. l. var. laticauda S. 1898 p. 287 pl. lxxii figs. 2,

A. nieszkowskii S. 1898 p. 287 pl. lxxii fig. 9, A. n. mut. jewensie pl. lxxii fig. 8, Lesnikova & Weber Atlas of the leading forms of the fossil faunas of the USSR Volume II 1949.

Asaphus angustifrons Dalman 1827 designated as type of Ptychopyge Angelin and placed on Official List of Specific Names p. 59, Opinion 538.

Asaphus hausmanni Brongniart 1822 placed on Official List of Specific Names and designated type of Odontochile Hawle & Corda 1847 p. 43, Opinion 537.

Asaphus powisi Murch. p. 235, Shackleton.

Asaphus ? sp. p. 1548, Bordet et al.

"Asperocare" argentinum Poulsen discussed and may belong to Chilometopus p. 8, Rusconi.

Asteropyge punctata (Steininger) subsp. p. 89 pl. figs. 1-6, Monseur.

Asteropyge? sp. p. 339 ff., Paulus.

Asthenaspis [misspelt Astenaspis on pl. expl.] gen. n. [Redlichiidae] p. 74 for type A. tenuis sp. n. p. 75 pl. vi figs. 10-16 text-figs. 5a-s Cambrian (Lower) U.S.S.R. (Yakutia), Suvorova Paleont. Zhurnal 1959 3.

Atractopyge aff. aspera (Linnarsson) p. 207ff., A. sp. pp. 206, 218, A. sp. ind. p. 214, **Dean** (1).

Australopyge see Thysanopyginae subfam. n.

Autoloxolichas reedi Phleger a synonym of Uralichas avus p. 102, Vaněk (2).

Basidechenella see Dechenella.

Basidechenella kayseri (R. Richter) p. 342, Paulus.

Basilicus kegelensis (Schmidt) 1898 p. 289 pl. lxxiv figs. 3-4, B. nobilis (Barrande) 1846 p. 289 pl. lxxiii figs. 4a-b, Lesnikova & Weber.

Basilicus tyrannus (Murchison) 1837 p. 30 pl. ii figs. 1-2, Balashova (1).

Bathynotidae see Bathynotina subord, n.

Bathynotina subord. n. [Redlichiida] p. 216 for Bathynotidae, Lochman-Balk Treatise on Invertebrate Paleontology Part O 1959.

Bathynotus lamanensis Lerm. p. 798, Z. A. Zhuravleva (1).

Bathyuriscellus see Jakutidae fam. n.

Bathyuriscina tannu-olae nov. gen. sp. nov. [nom. nud.] [Corynexochidae] p. 16, O. K. POLETAEVA in Pokrovskaya.

Bathyuriscus cf. adaeus text-fig. 3k, Kindle & Whittington.

 $Bathyuriscus \ ?$ sp. [det. O. K. Poletaeva] p. 16, Pokrovskaya.

Bathyurus ingalli Raymond, B. spiniger (Hall), B. trispinosus Wilson p. 85ff., B. aff. bandifer Sinclair p. 91, Colquboun.

Bathyurus sp. A, B. sp. B [both undescr.] p. 39, Thorsteinsson.

Bayfieldia text-fig. 3c, Kindle & Whittington.

Bayfieldia sp. undet. p. 388 pl. lv figs. 20-23, Rasetti.

Bellaspis ? oblonga sp. n. Cambrian (Upper) U.S.A. (Idaho) p. 413 pl. lvii figs. 23-24, Lochman & Hu J. Paleont. 33 3 1959.

Bellefontia sp. cf. B. chamberlaini Clark p. 113 pl. xxi figs. 18, 20, 22, Berg & Ross.

Bergeroniaspis Lermontova 1951 discussed with synonym Olekmaspis Suvorova and referred to Protoleninae p. 45, B. shanganus sp. n p. 46 pl. ii figs. 5-6, 8-10, 16, B. lebedevae sp. n, p. 50 pl. ii figs. 4, 7, B. sisovae sp. n, p. 52 pl. ii figs. 15, 17, B. cf. s. p. 32, B. argutus sp. n, p. 54 pl. ii figs. 3, 12, B. zaicevi sp. n, p. 56 pl. ii fig. 13 Cambrian (Lower) U.S.S.R. (Tuva), B. sp. p. 29, Pokrovskaya Trud. geql. Inst. S.S.S.R. 27 1959.

Bergeroniaspis Lermontova 1951 referred to Lermontoviidae; B. procera & B. nitens Suvorova excluded p. 395, **Sdzuy**.

Bergeroniaspis range in Altai-Sayan pl. i, Zhurav-leva & Repina.

Bergeroniaspis ornatus Lerm., B. divergens L. p. 1372, Z. A. Zhuravleva (2).

Bergeroniellus Lermontova 1940 referred to Lermontoviidae p. 395, **8dzuy.** 

Bergeroniellus divergens Lerm., B. kutorginorum L., B. ornatus L. p. 798, Z. A. Zhuravleva (1).

Bigotina Cobbold 1935 age considered probably lower L. Cambrian p. 402, Sdzuy.

Bigotininae Hupé 1952 [sic] discussed and referred to Dolerolenidae, taken to include Bigotina, Bigotinops, Pruvostina, Ouijjania, ? Pruvostinoides, ? Yumanocephalus p. 400, Sduy.

Bigotinops Hupé 1952 [sic] referred to Bigotininae p. 401, Sdzuy.

Billingsaspis adamsii formula of glabellar dimensions p. 481, Shaw.

Binodaspis range in Altai-Sayan pl. i, Zhuravleva & Repina.

Blackwelderia see Prodamesella gen. n.

Blackwelderia—list of spp. of Damesella & B. showing increase in no. of pygidial spines p. 233, B. octaepina [recte octoepina] Resser & Endo 1937 with synonym D. octaepina Kobayashi 1935 p. 233, Chang.

Boeckaspis kasachstanicus Bal. [not seen] p. 144, Balashova & Balashov.

Bohemopyge see Niobinae subfam. n.

Bolaspidella housensis (Walcott) p. 84, Bright.

Bolaspididae fam. n. [Norwoodiacea] p. 304 for Bolaspis, Acrocephalops, Eldoradia and Rawlinsella, Howell Treatise on Invertebrate Paleontology Part O 1959.

Bolaspis see Bolaspididae fam. n.

Bolaspis aemula sp. n. p. 204 pl. xxii fig. 15 pars, pl. xxiii figs. 1-2, B. f sp. undet. p. 204 pl. xix fig. 4

pars Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Bonnia Walcott discussed, B. sisovae sp. n. p. 135 pl. x figs. 5-6, B. asiatica sp. n. p. 137 pl. x figs. 4, 11, 14, B. dubia [misspelt bubia on pl. expl.] sp. n. p. 140 pl. x fig. 8 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Bonniarange in Altai-Sayan pl. i, Zhuravleva & Repina.

Bonnia sp. p. 1686, Theokritoff.

Bonnia sp. p. 1032, Zhuravleva et al. (2).

Borogothus see Promegalaspides.

Borogothus see Promegalaspidinae subfam. n.

Bowmania pennsylvanica sp. n. p. 395 pl. lv figs.1-5 text-fig. 1 Cambrian (Upper) U.S.A. (Pa, ? Md); reference of Anomocare parvula Weller to B. unconvincing p. 396, Rasetti J. Paleont. 33 3 1959.

Briscoia celtica (Salter) horizon discussed p. 222, Shackleton.

Broeggerolithus harnagensis (Bancroft) p. 233 ff., B. cf. h. p. 249, B. soudleyensis (B.), B. broeggeri B., B. sp. p. 235, Shackleton.

Broeggerolithus nicholsoni (Reed), B. aff. n. p. 214, B. cf. n. p. 206, B. transiens (Bancroft), B. sp. p. 207, **Dean (1).** 

Brongniartella see Homalonotus.

Brongniartella bisulcata (Salter) p. 235ff., Shackle-ton.

Brongniartella cf. parva Harper, B. cf. rudis (M'Coy) p. 206, B. bisulcata (M'Coy), B. sp. ind. (small) p. 207, B. cf. b., B. sp. ind. p. 214, B. sp. nov., B. sp. (? nov.) [both undescr.] p. 208, B. ascripta (Reed) p. 214, Dean (1).

Bronteidae see Thysanopeltidae.

Bronteopsis scotica Etheridge et Nicholson pl. v figs. 13-16, B. nannus sp. nov. pl. v figs. 17-18, Balashova (1).

Bronteopsis scotica Nicholson et Etheridge p. 29 [pl. v figs. 13–16], B. nannus sp. n. p. 32 [pl. v figs. 17–18] Ordovician (Middle) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Bronteopsis sola (Barr.) for Ogygia s. B. 1872 p. 215, Vaněk (1).

Bronteus aculeatus Weber type of Weberopeltis gen. n. [quod vide].

Bronteus romanovskii Weber 1948 p. 298 pl. lxxix figs. 2a-b, B. aculeatus W. 1945 p. 298 pl. lxxix figs. 1a-b, Lesnikova & Weber.

Bulaiaspis peleduicus Repina, B. vologdini Lerm., B. primus L. p. 797, Z. A. Zhuravleva (1).

Bulaiaspis tasevicus Rep., B. primus Lerm., B. vologdini L., B. manicus R. [nom. nud.] p. 1372, Z. A. Zhuravleva (2).

Bullaglabellaspis Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Bun pl. lx

[195

Bun

p. 297 pl. lxx pl. iii 23–29

Balas

But Ordov sp. n. p. 48 (Arcti [pls. i

> Ca Ca Ca

Ru

Calla Inve Ca Ca

P. 30 P. 30 Bala

1a, 16, Ho

p. d Ba

> Sileste p.

pl

p

15

Ó

Bumastides bedpakensis Weber 1948 p. 291 pl. lxxxi figs. 1-4, Lesnikova & Weber.

Bumastus barriensis Murchison 1839 p. 297 pl. lxxviii figs. 7a-b, B. nudus (Angelin 1854) p. 297 pl. lxxviii fig. 8, B. milleri Billings 1859 p. 297 pl. lxxviii figs. 9a-b, Lesnikova & Weber.

Bumastus holei Foerste var. taimyrica var. nov. pl. iii figs. 20-22, B. taimyricus sp. nov. pl. iii figs. 23-29, B. nordicus sp. nov. pl. iii figs. 30a-r, Balashova (1).

Bumastus holei Foerste var. taimyrica var. n. Ordovician (Upper) p. 45 [pl. iii figs. 20-22], B. nordicus sp. n. p. 46 [pl. iii figs. 30a-r], B. taimyricus sp. n. p. 48 [pl. iii figs. 23-29] Silurian (Lower) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Bumastus sp. A, B. sp. B p. 36, B. sp. C [all undescr.] p. 37, Thorsteinsson.

Calipernurus insolitus p. 385, Whittington.

Callavia see Callaviinae subfam. n.

Callaviinae subtam. n. [Olenellidae] p. 192 for Callavia, Judomia, Kjerulfia, Poulsen Treatise on Invertebrate Paleontology Part O 1959.

Calliops narrawayi Okulitch p. 91, Colquhoun.

Calliops sp. cf. C. narrawayi Okulitch p. 39, Thorsteinsson.

Calliops strasburgensis, C. callirachis p. 385, C. sp. p. 386, C. loxorachis p. 387, Whittington.

Calliops taimyricus sp. nov. pl. v figs. 9-12, Balashova (1).

Calodiscus lobatus p. 1686, Theokritoff.

Calymene ex gr. blumenbachi Brongn. p. 594 text-fig. la, C. macrocephala Maksimova [nom. nud.] text-fig. le, C. sp. text-fig. le, Maksimova & Organova.

Calymene neointermedia R. & E. Richter p. 1039, Holland et al.

Calymene neointermedia R. & E. Richter, C. sp. p. 514, Walmsley.

Calymene taimyrica sp. nov. pl. i figs. 11-14, Balashova (1).

Calymene taimyrica sp. n. p. 52 [pl. i figs. 11-14] Silurian (Lower) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Calymene tenera Barrande referred to Platycalymene p. 215, Vaněk (1).

Calymene tuberculata Bruennich 1885 [sic] p. 324 pl. xxvii figs. 7-8, Strnad (8).

Calymene tuberculata (Brünnich 1781) p. 306 pl. lxxxii fig. 9, Lesnikova & Weber.

Calymene sp. p. 477, Holland.

Calymene sp. p. 235 ff., "C." sp. p. 233, Shackleton.

Calymene sp. A [undescr.], C. sp. p. 50, Thorsteinsson.

Calymenella (Eohomalonotus) bohemicus (Barr.) p. 48, Vaněk (3). Calymenid pygidium, Calymenid pygidium ind. p. 207. Dean (1).

Calyptaulaz (Subgen. A) for Homalops Remelé 1885 (non Motschulsky 1850) p. 492, STRUVE in Harrington et al.

Calyptaulax calderi Wilson p. 91, C. aff. c. p. 88, Colquhoun,

Calyptaulax aff. norvegicus Størmer p. 207ff., Dean (1).

Caphyra taimyrica sp. n. p. 28 pl. i fig. 7, C. ? kluevkaensis sp. n. p. 28 pl. i fig. 1 Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 14 1959.

Carinopyge spinifera sp. nov. pl. iv figs. 18a-r, Balashova (1).

Catadoxides Matthew 1899 discussed and considered not a Dolerolenid p. 398, Sdzuy,

Cedarina prima Lochman 1944 & C. cordillerae (Howell & Duncan 1939) considered ♂ & ♀ of one sp. p. 415, Lochman in Lochman & Hu.

Centropleura sp. p. 359, Miroshnikov et al.

Ceratarges 800 Lichas.

Ceratarges sp. p. 349, Paulus.

Ceratarginae Tripp 1957 discussed and taken to include Hemiarges, Perunaspis, Lobopyge, Acanthopyge, Eifliarges, Radiolichas, Ceratarges, Craspedarges, Acantharges p. 118 [also Trochurus p. 144]; Dicranogmus referred to Lichainae p. 116, Vaněk (2).

Ceratevenkaspis pentagonus sp. nov. pl. iv figs. 7-8, C. taimyricus sp. nov. pl. iv figs. 9-11, 13a-6, C. ? borealicus sp. nov. pl. iv figs. 12a-6, C. parnaicus sp. nov. pl. iv fig. 14, Balashova (1).

Ceratocephala triacantheis p. 384, C. laciniata, C. (Ceratocephalina) tridens p. 385ff., C. rarispina p. 387, Whittington.

Ceratocephala vesiculosa Beyrich [for Ferronnière 1920-22 pl. 5 fig. 100] p. 103, Pillet (2).

Ceratocephalina soo Ceratocephala,

Ceratolithus cornutus n. gen. n. sp. [nom. nud.] p. 43 Ordovician China, Lu.

Ceratopyge forficula Sars 1847 p. 274 pl. lxv figs. 4a-c, Lesnikova & Weber.

Ceratopyge forficula (Sars) p. 274, Tjernvik.

Ceratopyge forficula f. (Sars), C. f. acicularis (Sars & Boeck) listed p. 174, **Henningsmoen**,

Ceratopyge cf. forficula Sars p. 144, Balashova & Balashov.

Ceratopyge transversa n. sp. [nom. nud.] p. 88 Ordovician China, Lu.

Ceraurinella chondra p. 384 ff., C. typa p. 385 ff., Whittington.

Ceraurinus borealicus sp. nov. pl. iii fię: 31, pl. v fig. 22, C. confluens Barton pl. v fig. 1, C. nordicus sp. nov. pl. iv fig. 19, pl. v fig. 2, C. taimyricus sp.

nov. pl. v figs. 3-4, C. marginatus B. pl. v fig. 5, C. aff. tenuisculptus Bradley pl. v fig. 6, C. kluevkaensis sp. nov. pl. v fig. 7, C. icarus (Billings) pl. v figs. 23 a-6, Balashova (1).

Ceraurinus daedalus Cox p. 37, Thorsteinsson.

Ceraurus dentatus Raymond & Barton, C. aff. pleurexanthemus Green p. 85, Colquhoun.

Ceraurus exsul (Beyrich) 1846 p. 307 pl. lxxxiii figs. 9–10, C. e. var. macrophtalmus Kutorga 1854 p. 308 pl. lxxxiii fig. 16, Lesnikova & Weber.

Ceraurus horridus Troedsson, C. sp. p. 37, Thorsteinsson.

Ceraurus—comparison of hypostome with that of Remopleurides p. 395 text-figs. 3a-B [C. sp.], C. whittingtoni p. 388, C. sp. p. 387, Whittington.

Chancia sp. p. 84, Bright.

Chasmops cf. extensa (Boeck) p. 207, C. aff. maxima (Schmidt) p. 208, C. sp. p. 214, C. cf. marri (Reed) p. 218, Dean (1).

Chasmops inge Rõõm. pl. xxv fig. 3, C. wesenbergensis Schm. pl. xxv fig. 5 pars 4, Kaljo (1).

Chasmops odini (Eichwald) 1840 p. 313 pl. lxxxv figs. 9a-b, C. o. var. itferensis Schmidt 1881 p. 313 pl. lxxxv fig. 10, C. marginata S. 1881 p. 313 pl. lxxxv figs. 7a-8, C. bucculenta Sjögren 1852 p. 314 pl. lxxxiv figs. 11a-b, C. wesenbergensis S. 1881 p. 314 pl. lxxxiii figs. 7-8, pl. lxxxv figs. 4-5, C. wrangeli S. 1881 p. 314 pl. lxx figs. 3a-b, pl. lxxxv figs. 6a-b, C. maxima S. 1881 p. 315 pl. lxxxv figs. 1-3, Lesnikova & Weber.

Chasmops odini (Eichwald 1840) from glacial boulder p. 130 text-fig., Strnad (3).

Chasmops odini (Eichwald 1840) p. 320 pl. xxvi fig. 5, pl. xxvii fig. 4, Strnad (8).

Chasmops of. wesenbergensis Schm. p. 6, C. sp. p. 7, C. eichwaldi S. p. 8ff., Männil.

Chasmops sp. p. 240, Shackleton.

Cheirurids p. 385, Whittington.

Cheiruroides Kobayashi discussed p. 124, C. maslovi sp. n. p. 126 pl. ix figs. 10-12 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.R. 27 1959.

Cheirurus sinicus <br/>n. sp. [nom. nud.] p. 77 Ordovician China, Lu.

Chelidonocephalus King 1937 discussed p. 223, Chang.

Chilometopus parabolicus Rusc. discussed; Asperocare argentinum Poulsen may belong to C. p. 8, Rusconi.

Chomatopyge marginifera, C. falcata p. 384, Whittington.

Chondragraulina Pokrovskaya [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Chondragraulis [sic] minussinensis Lerm. p. 626, Anatol'eva & Shelkovnikov.

Chondragraulos range in Altai-Sayan pl. i, Zhurav-leva & Repina.

Chosenia divergens n. sp. [nom. nud.] p. 34 Ordovician (Lower) China, Lu.

Chungkingaspis plethopeltoides n. gen. n. sp. [nom. nud.] p. 36 Ordovician (Lower) China, Lu.

Clavaspidella kanabensis sp. n. p. 195 pl. xix figs. 29-32, C. sp. undet. p. 196 pl. xxii fig. 16 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Coignouina see Otarion.

Collis gen. n. [Trinucleidae] for type C. parvulus sp. n. p. 1313 pl. figs. 2a, b, e, text-fig. 4b, C. sp. pl. figs. 2z, b, e Ordovician (Upper) U.S.S.R. (Kazakhstan), Koroleva C. R. Acad. Sci. U.R.S.S. (ns) 124 6 1959.

Conococheaguea gen. n. [affinities obscure] for type C. ovata sp. n. p. 396 pl. lii figs. 19-26 text-fig. 2 Cambrian (Upper) U.S.A. (Pa & Md), Rasetti J. Paleont. 33 3 1959.

Conocoryphe sulzeri Schlot. p. 110 (pl. 36) fig., C. s., var. latus and var. altus p. 110, C. ribeiro Barr. & Vern. p. 111 (pl. 37) fig., Badillo.

Conokephalina abdita (Salter) horizon discussed p. 222, Shackleton.

Conokephalina termieri & C. tienfongensis Mansuy spp. belong to new subgenus of Proasaphiscus p. 227 n, Chang.

Conolichas eichwaldi (Nieszk.), C. ef. e. p. 7, Männil.

Conolichas melmerbiensis (Reed) p. 214, Dean (1).

Coosia sp. p. 359, Miroshnikov et al.

Cornuproetus (C.) buchi taciturnus Kegel [for Ferronnière 1920–22 pl. 1 figs. 11, 12] p. 102, C. (C.) b. platykraspedon Erben p. 103, Pillet (2).

Coronocephalus Grabau 1924 emend. diagnosis p. 80, distribution p. 76, C. kobayashii sp. n. for Encrinurus sp. Hamada 1958 Silurian (Middle) Japan (Kyushu) p. 80 pl. vi figs. 1–18 text-fig. 2 (reconstrs.), C. kitakamiensis (Sugiyama) 1941 discussed p. 84 pl. vi figs. 19–20, Hamada Japan. J. Geol. Geogr. 30 7 1959.

Corrugatagnostus convergens sp. n. including Olenus? of Baily 1862 partim p. 370 pl. lxii figs. 17-19 Ordovician (Upper) Eire, Weir Palaeontology 1 4 1959.

Corrugatagnostus transitans n. sp. [nom. nud.] p. 77 Ordovician China, Lu.

Corydocephalus flabellatus pl. x fig. 1, C. interjectus pl. x fig. 4, C. propinquus pl. xi fig. 3, C. verrucosus pl. xi fig. 2 Hawle & Corda spp. lectotypes chosen and figured and all synonyms of Trochurus epeciosus p. 147, Vaněk (2).

Corynexochidae see Bathyuriscina, Eniseja, Tuvinia.

Corynexochoidea see Jakutidae fam. n.

Crepicephalina damia var. rectangula var. n. p. 209 (232) pl. iv figs. 3-4 Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Ferro p. 103

[198

Cro

a dist

Cry (incluprotafigs. I p. 38'

> Cte (pl. 3

figs. 8
Cy
Cybe

Ham

Cy Bala Cy Chin

Cy 4a-b

Consimply Van

figs.
figs.
C
Ric

fron

P. 3

I Tri Mo:

Sil

Cromus Barrande 1850 [sic] discussed and considered a distinct genus p. 76, Hamada.

Crotalocephalus sternbergi Boeck p. 102 [for Ferronnière 1920-22 pl. 1 fig. 5], C. gibbus Beyrich p. 103, Pillet (2).

Cryphops see Phacops.

Cryptolithids p. 233ff, Shackleton.

Cryptolithus tesselatus Green 1832 morphology (including hypostome) and ontogeny (including protaspis) described p. 441 ff. pl. xx figs. 1-11, pl. xxi figs. 1-19, pl. xxii figs. 1-12, pl. xxiii figs. 1-23, C. sp. p. 387, Whittington.

Cryptolithus sp. p. 240, Shackleton.

Ctenocephalus coronaatus Barr. [var. altus] p. 112 (pl. 38) fig., Badillo.

Cybele bellatula Dalman 1826 p. 307 pl. lxxxii figs. 10a–11b, C. planifrons Weber 1948 p. 307 pl. lxxxii fig. 8, Lesnikova & Weber.

Cybelinae Holliday 1942 emend. taken to include Cybele, Atractopyge, Cybeloides, Ectenonotus, Cybellela, Cybelopsis, Bevanopsis and Paracybeloides p. 73, Hamada.

 $Cyclognathus\ micropygus\ Linn.\ p.\ 144,\ Balashova\ \&\ Balashov.$ 

Cyclopyge radiata n. sp. [nom. nud.] p. 77 Ordovician China, Lu.

Cyclopyge sp. p. 223, Shackleton.

"Cyclopyge" sp. p. 46ff., Vaněk (3).

Cyclopygid gen. et sp. indet. p. 379 pl. lxiii figs. 4a-b, Weir.

Cyphaspis (Schmidtella) uralica Tschernyschew 1893 p. 300 pl. lxxix figs. 9a-b, Lesnikova & Weber.

Cyphinium see Griffithides.

Cyphoproetus depressus (Barrande) for Dicranopeltis simplex of Hawle & Corda 1847 non B. 1846 p. 117, Vaněk (2).

Cyrtometopus affinis Angelin 1854 p. 310 pl. lxxxiii figs. 4–6, C. clavifrons (Dalman) 1826 p. 310 pl. lxxxiii figs. 13–15, Lesnikova & Weber.

Cyrtosymbole (Waribole) cf. warsteinensis R. & E. Richter, C. (W.) avita (R. & E. R.) p. 104, C. (W.) a. from Carboniferous p. 78, Lewowicki,

Cyrtosymbole sp. p. 266 pl. i fig. 3, Pfeiffer.

Dactylocephalus breviceps, D. hsūi p. 33, D. laevigata p. 34, D. obesus, D. ? abnormis p. 38 n. spp. [nom. nuda] Ordovician China, Lu.

Dalmanitacea see Phacopina subord, n.

Dalmanites Barrande 1852 designated type of class Trilobita Walch 1771 p. 172, HARRINGTON, HENNINGS-MOEN & MOORE in Harrington et al.

Dalmanites see Phacops.

Dalmanites batalleri sp. n. p. 5 [6 figs. on second pl.] Silurian (Lower) Spain (Gerona), Sampelayo Notas Inst. geol. Esp. No. 13 1944. Dalmanites batallieri Sampelayo p. 25, D. sp. p. 26, Suñer Coma & Fernández de Villalta.

Dalmanites vulgaris (Salter) p. 1038, Holland et al.

Dalmanites vulgaris (Salter) p. 514, Walmsley.

Dalmanitids p. 384ff., Whittington.

Dalmanitina nanchengensis Lu p. 78, Lu.

Dalmanitina proaeva proaeva (Emmrich), D. p. čilinensis Šnajdr, D. p. socialis (Barr.) p. 47, Vaněk (3).

Dalmanitina proaeva ? Barr. p. 1548, Bordet et al.

Dalmanitina sp. nov. [undescr.] p. 208, Dean (1).

Damesella Walcott 1905 discussed with list of spp. of D. & Blackwelderia showing increase in no. of pygidial spines, D. bilongispina sp. n. p. 209 (232) pl. iv fig. 5 Cambrian (Middle) China (Shantung); D. octaspina Kobayashi 1935 a synonym of B. octaspina Resser & Endo 1937 p. 233, D. quadrata [Resser &] Endo referred tentatively to Prodamesella gen. n. p. 219, Chang Acta palaeont. sinica 7 3 1959.

Damesella soo Prodamesella gen. n.

Dechenella (Basidechenella) dombroviensis (Gürich 1896) p. 136, Strnad (4).

Dechenella sp. p. 740, McLaren.

Deiphoninae see Ovalocephalus gen. n.

Denckmannites see Phacopidella.

Despujolsia see Despujolsiidae fam. n.

Despujolsiidae fam. n. [Redlichiacea] for Despujolsia p. 204, Harrington Treatise on Invertebrate Paleontology Part O 1959.

Diacalymene marginata Shirley p. 218, D. cf. m. p. 208, Dean (1).

Diacanthaspis lepidus, D. secretus, D. ulrichi p. 385ff., D. orandensis, D. scitulus, D. aff. u. p. 387, D. cooperi p. 388, Whittington.

Dianops see Phacops.

Diaphanometopinae subfam, n. [Pliomeridae] p. 445 for *Diaphanometopus*, Jaanusson Treatise on Invertebrate Paleontology Part O 1959.

Diaphanometopus see Diaphanometopinae subfam.

Dicranogmus Hawle et Corda 1847 discussed and referred to Lichainae p. 116, D. simplex (Barrande 1846) excluding record of H. & C. [to Cyphoproctus depressus] and with synonym D. pustulatus H. & C. of which lectotype chosen and figured p. 117 pl. i figs. 4-7, Vanék (2).

Dicranopeltis Hawle & Corda 1847 discussed p. 108, D. scabra scabra (Beyrich 1845) emend. Prantl & Vaněk 1958 with synonym D. aspera H. & C. of which lectotype chosen and figured p. 109 pl. iv figs. 1, 6-9, pl. xii fig. 1, text-figs. 10-12, D. s. propinqua (Barrande 1846) emend. P. & V. with synonym i.a. D. granulosa H. & C. of which lectotype chosen and figured p. 112 pl. iv figs. 5, 10, pl. xii fig. 2, text-fig. 13, D. scabra inc. subspec. pl. iv figs. 2-4; D. † balkhaschicus and D. † turensis Weber 1951 tentatively synonyms of Trochurus speciosus p. 145, Vaněk (2).

Dienstina soo Phacopidella.

Digonus see Homalonotus.

'Dikellocephalus' celticus Salter horizon p. 56, Stubblefield (1),

Dikelocephalid free cheek p. 222, Shackleton.

Dikelocephalidae see Olenina.

Dikelocephalina gibba n. sp. [nom. nud.] p. 76 Ordovician China, Lu.

Dikelocephaloidea — Granulariidae referred, relations to Asaphoidea discussed p. 106, Suvorova (1).

Dikelokephalina dicraeura (Angelin) p. 144, Balashova & Balashov.

Dikelokephalina dicraeura (Angelin 1854) p. 163 pl. i figs. 1-4, Henningsmoen.

Dikelokephalininae discussed and referred to Hungaiidae p. 161, Henningsmoen.

Dimeropyge spinifera, D. virginiensis p. 384, D. sp. p. 387, Whittington.

Dindymene ovalis sp. n. including D. haidingeri of Baily 1862 p. 380 pl. lxii figs. 9-10 Ordovician (Upper) Eire, Weir Palaeontology 1 4 1959.

Dindymene sp., D. sp. nov. [undeser.] p. 207, **Dean** (1).

Dindymeninae subfam. n. [recte Přibyl 1953] p. 448, Henningsmoen in Harrington et al.

Dinesidae see Erbina and Sayanella genn. n.

Dionide aff. richardsoni Reed p. 372 pl. lxiii figs. 7–8, Weir.

Ditomopyge from W. Australia p. 67ff., McWhae et al.

Dokimocephalidae see Minkella gen, n.

Dolerolenidae Kobayashi 1935 discussed p. 393ff., classified into Doleroleninae and Bigotininae p. 390, Sdzuy.

Doleroleninae Kobayashi 1935 discussed with synonyms Metadoxidinae p. 396 and Abadiellidae p. 398, taken to include Dolerolenus, Metadoxides, Abadiella, Lunolenus nom. nud., ? Redlichina, ? Wutingaspis p. 390, Sdzuy.

Dolerolenus Leanza 1949 discussed p. 396, D. formosus n. sp. (nom. nud.) p. 403 pl. i fig. 5 (reconstr.), D. n. sp. [undescr.] pl. i fig. l, **Sdzuy.** 

Dolichoharpes reticulata p. 384ff., Whittington.

Dolichometopinae see Erbiopsidella and Rondocephalus genn, n.

Dorypyge sp. indet. p. 359, Miroshnikov et al.

Drevermannia (D.) richteri sp. n. for D. sp. A Pfeiffer 1954 p. 267 pl. ii figs. 1-4, D. sp. p. 268 pl. ii figs. 5-6 Devonian (Upper) Germany (Thuringia), Pfeiffer Geologie 8 3 1959.

Drumaspis walcotti Resser with synonyms idahoensis, briscoensis & nitida R. 1942 p. 416 pl. lx figs. 1-14, Lochman & Hu.

Duftonia gen. n. [Acastinae] p. 143 for type D. lacunosa sp. n. including probably Pterygometopus sp. Bancroft 1948 p. 144 pl. xix figs. 1-3, 5, 6, 8 Ordovician (Upper) Britain (N. England), D. aff. lacunosa p. 147 pl. xix figs. 4, 7, 9-11, Dean Palaeontology 2 1 1959.

Duftonia lacunosa Dean p. 208, Dean (1).

Dunderbergia text-fig. 3h, Kindle & Whittington.

Dytremacephalus text-fig. 3g, Kindle & Whittington.

Eccoptochile (E.) clavigera (Beyrich), E. (Eccoptochiloides) tumescens (Barr.) p. 47, Vaněk (3).

Eccoptochiloides see Eccoptochile.

Ectillaeninae subfam. n. [Illaenidae] p. 374 for Ectillaenius, Zbirovia and Zdicella, Jaanusson Treatise on Invertebrate Paleontology Part O 1959.

Ectillaenus see Ectillaeninae subfam, n.

Edelsteinaspidae see Paleofossus gen. n.

Edelsteinaspididae see Nodiceps gen. n.

Edelsteinaspis see Laticephalus gen. n.

Edelsteinaspisrange in Altai-Sayan pl. i, Zhuravleva & Repina.

Ehmaniella arizonaensis sp. n. p. 205 pl. xxiii fig. 4, E. basilica sp. n. p. 206 pl. xx figs. 19–20 partes, pl. xxiii figs. 10–13, E. hebes sp. n. p. 206 pl. xxiv figs. 2–6 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg, Instn No. 563 1945.

Eldoradia see Bolaspididae fam. n.

Elganellus range in Altai-Sayan pl. i, Zhuravleva & Repina.

Eleganolimba gen. n. [Aldonaidae] p. 97 for type E. fibrata 5p. n. p. 98 pl. ix fig. 13 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.R. 27 1959.

Elegestina gen. n. [Protoleninae] p. 63 for type E. antiqua sp. n. p. 64 pl. i fig. 11 Cambrian (Lower) U.S.S.R. (Tuva), E. sp. p. 29, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Elganellus acceptus Suv., E. probus Suv. p. 797, Z. A. Zhuravleva (1).

Elganellus acceptus Suv., E. probus S., E. zhuravlevae sp. n. [? Pokrovskaya] [nom. nud.] p. 1372, Z. A. Zhuravleva (2).

Ellipsocephalacea see Redlichiina subord. n.

Ellipsocephalidae (indet.) p. 771, Orłowski (1).

Ellipsocephaloides butleri Resser p. 111 pl. xxii fig. 8. Berg & Ross.

Ellipsocephalus polytomus Lnrs., E. (other spp.) p. 771, Orlowski (1).

Ellipsocephalus polytomus Linnarsson p. 515 pl. i figs. 1-3, 5, 9 [non 4], E. gürichi sp. n. p. 516 pl. i figs. 6-8, 10, 4 [non 9], text-fig. 1a, E. puschi sp. n. p. 517 pl. ii figs. 1a-c, text-fig. 1b, E. sandomiri sp. n. p. 518 pl. ii figs. 2-4, text-fig. 1c, E. jugoszovi sp. n. p. 518 pl. ii figs. 5-6, text-fig. 1d, E. longus sp. n. p. 519 pl. ii figs. 7-9, text-fig. 1e Cambrian (Middle) Poland (H. Cross Mts.), Orlowski Bull. Acad. polon. Sci. [sci. chim., etc.] 7 7 1959.

Gern El pres

[19

El

popu 5–7 E pl. x Res

E. (Mi Ins

int

Fa

of property of the contract of

1

Ellipsocephalus sancta-crucensis Czarnocki see Germaropyge.

Elliptocephala asaphoides fauna p. 1686, Theokritoff.

Elrathia kingii (Meek) discussion of morphology, preservation, habits and biometries of a topotype population p. 85ff. pl. xvii figs. 1-10, pl. xviii figs. 1-2, 5-7 (sections), E. k. l p. 88, Bright.

Elrathia nitens sp. n. p. 209 pl. xxvii fig. 35, E. sp. pl. xxi fig. 16 pars Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Elrathiella decora sp. n. p. 209 pl. xxiv fig. 1, E. f ineueta sp. n. p. 210 pl. xxv figs. 5-8 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Elrathina cordillerae (Rominger) p. 84, Bright.

Elviniidae see Olenina.

Encrinuridae Angelin 1854 placed on Official List of Family-Group Names p. 44, **Opinion 537.** 

Encrinuridae Angelin 1854 discussed and classified into Encrinurinae, Cybelinae, Dindymeninae and Staurocephalinae p. 72ff., Hamada.

Enerinurids p. 384ff., Whittington.

Encrinurinae discussed and classified into Encrinuroides, Encrinurus multisegmentatus group, Coronocephalus, Encrinurus konghsaensis group, Encrinurus, Cromus and Mitchellia p. 74 ff., Hamada.

Encrinurinae see Mitchellaspis nom. n.

Encrinurus Emmrich 1844 placed on Official List of Generio Names, designation of Entomostracites punctatus Wahlenberg [quod vide] as type p. 43, Opinion 537.

Encrinurus—several Asian records discussed p. 78ff., E. spp. Kayser 1883 discussed and casts refigured p. 84 pl. vi figs. 21–22 [see also Coronocephalus and Cromus], Hamada.

Encrinurus punctatus (Wahlenberg), E. sp. p. 514, Walmsley.

Encrinurus punctatus (Wahl.) pl. xxiv figs. 1-2, Kalio (1).

Encrinurus seebachi Schmidt 1881 p. 306 pl. lxxxii figs. 4-5 E. punctatus (Wahlenberg) 1821 p. 306 pl. lxxxii figs. 6a-7b, Lesnikova & Weber.

Encrinurus sp. p. 214, Dean (1).

Encrinurus sp. p. 1039, Holland et al.

Encrinurus sp., E. sp. ? p. 7, Männil.

Encrinurus sp. p. 49ff., E. n. sp. A [undescr.], E. sp. of, E. moderatus Poulsen, E. n. sp. [undescr.] p. 50, Thorsteinsson.

Eniseja tannu-olae nov. gen. sp. nov. [nom. nud.] [Corynexochidae] p. 16, O. K. POLETAEVA in Pokrovskaya.

Entomaspididae see Harpina subord. n.

Entomaspis radiatus Ulrich p. 393 pl. lii figs. 9, 10 [pygidium tentatively assigned]; E. recorded by A. R. Palmer from Nevada p. 377, Rasetti.

Vol. 96

Entomostracites punctatus Wahlenberg 1821 validated (by suppression of Trilobus p. Brünnich 1781) and placed on Official List of Specific Names and designated type of Encrinurus Emmrich 1844 p. 43, lectotype figured pl. i figs. 1–2, Opinion 537.

Eoacidaspididae see Odontopleurida ord. n.

Eocorycephalus see Synphoria,

Eodiscidae discussed and classified into Eodiscinae and Weymouthiinae p. 163, Pokrovskaya.

Eodiscinae see Ladadiscus gen. n.

Eodiscus see Microdiscus.

Eodiscus scanicus text-fig. 3j, Kindle & Whittington.

Echarpes sp. indet. p. 91, Colquhoun.

Eohomalonotus see Calymenella.

Eophacops tentatively recorded p. 41, Williams.

Erbia range in Altai-Sayan pl. i, Zhuravleva & Repina.

Erbia granulosa Lerm. p. 1033, Zhuravleva et al. (2).

Erbia ex gr. sibirica Lerm. p. 626, Anatol'eva & Shelkovnikov.

Erbina gen. n. [Dinesidae] p. 148 for type E, rara sp. n. p. 149 pl. vii figs. 7, 10 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Erbiopsidella gen. n. [Dolichometopinae] p. 115 for type E. convexa sp. n. p. 117 pl. vi figs. 1-9 Cambrian (Lower) U.S.S.R. (Tuva), E. (young form) pl. x fig. 16 pars, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Erbiopsidella range in Altai-Sayan pl. i, Zhuravleva & Repina.

Erbiopsis Lermontova discussed p. 119, E. grandis L. lectotype selected [illegitimate] p. 120 pl. vi fig. 12, Pokrovskaya.

Erbiopsis range in Altai-Sayan pl. i, Zhuravleva & Repina.

Eremiproetus eremita erbeni Pillet p. 103, Pillet (2). Estoniops exilis (Eichw.) pl. xxix figs. 2-3, Kaljo (2).

Estoniops jukesi (Salter) so referred p. 146, Dean (2).

Estoniops jukesi (Salter) p. 207, Dean (1).

Euarges sp. p. 109, Thorsteinsson.

Eucalymene quadratus n. gen. n. sp. [nom. nud.] p. 46 Ordovician China, Lu.

Eudolatites angelini (Barr.) p. 48, Vaněk (3).

Euloma (possibly) p. 223, Shackleton.

Euloma ornatum Angelin listed p. 174, Hennings-meen.

Euloma ornatum Angelin p. 274, Tjernvik.

Euloma ornatum Ang. var. alimbetica Bal., E. kasachstanica B., E. kelleri B. [none seen] p. 144, Balashova & Balashov. Euptychaspis typicalis Ulrich p. 393 pl. lii figs. 11–13,

Evenkaspis norilskiensis sp. nov. pl. iv figs. 15-17, Balashova (1).

Eymekops rectangularis Resser & Endo [recte Endo] type of Peichiashania gen. n. [quod vide] p. 223, Chang.

Falanaspis aliena Tjernvik 1956 p. 171 pl. i fig. 8, Henningsmoen.

Flexicalymene incerta (Barr.) p. 48, Vaněk (3).

Flexicalymene onniensis Shirley, F. cf. laticeps Bancroft p. 207, F. aff. o. p. 208, F. cf. caractaci (Salter), F. aff. c. p. 214, **Dean** (1).

Flexicalymene senaria p. 388, F. sp. p. 387, Whittington.

Fordaspis nana p. 1686, Theokritoff.

Fuchouia spinosa sp. n. p. 210 (235) pl. iv figs. 12–16 text-figs. 26–27 Cambrian (Middle) China (Shantung); F. quadrata Resser & Endo attributed free cheek R. & E. 1937 pl. 39 fig. 4 pars) referred to Szesspis sp. p. 230, Chang Acta palaeont. sinica 7 3 1959.

Geragnostus crassus Tjernvik, G. sidenbladhi (Linnarsson) from Norway p. 174, Henningsmoen.

Geragnostus sidenbladhi Linn. p. 144, Balashova & Ralashov.

Geragnostus yangtzeensis n. sp. [nom. nud.] p. 33 Ordovician (Lower) China, Lu.

Geragnostus sp. undet. p. 380 pl. li figs. 15–18, Rasetti.

Germaropyge sancta-crucensis (Czarnocki) [sp. n.] neotype [sic] for Ellipsocephalus s.-c. C. 1927 nom. nud. p. 527 pl. ii figs. 4-12 Cambrian (Lower)-Poland (H. Cross Mts.), Samsonowicz Bull. Acad. polon. Sci. [sci. chim., etc.] 7 7 1959.

Gigantopygidae fam. n. [Redlichiacea] for Gigantopygus p. 204, Harrington Treatise on Invertebrate Paleontology Part O 1959.

Gigantopygus see Gigantopygidae fam. n.

Glaphurina strigata Weber 1932 p. 305 pl. lxxix figs. 7a-b, G. aff. lamettensis [sio] Ulrich 1929 p. 305 pl. lxxix fig. 8, Lesnikova & Weber.

Glaphurus altaicus Weber 1948 p. 305 pl. lxxix fig. 10, Lesnikova & Weber.

Glossopleura in Mendoza discussed p. 7, Rusconi.

Glossopleura mckeei Resser lectotype selected and figured p. 196 pl. xxi figs. 1–5, G. tuta sp. n. p. 196 pl. xxvi figs. 5–6, G. sp. undet p. 197 pl. xx figs. 21–22, pl. xxi fig. 6 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Glyphaspis robustus Deiss p. 1781, Kauffman.

Glyphaspis vulsa sp. n. p. 198 pl. xxii figs. 12–14, G. tecta sp. n. p. 198 pl. xxii fig. 15 pars, G. kwaguntensis sp. n. p. 199 pl. xxv figs. 12–13 partes, pl. xxvi figs. 2–4 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Glyptagnostus [reticulatus p. 10] text-fig. 3f, Kindle & Whittington.

Glyptagnostus reticulatus (Ang.) p. 360, Miroshnikov

Glyptometopus laflammei (Clark) p. 381 pl. li figs. 30-31, Rasetti.

Goldillaeninae subfam. n. [Scutelluidae] p. 34 for Goldillaenus, Goldillaenoides gen. n. and Illaenus indeterminatus Walcott, Balashova Sborn. stat. paleont. biostrat. 15 1959.

Goldillaenoides taimyricus gen. et sp. nov. pl. i figs. 8-9, pl. iii figs. 32a-6, Balashova (1).

Goldillaenoides gen. n. [Goldillaeninae] p. 35 for type G. taimyricus sp. n. p. 36 [pl. i figs. 8-9, pl. iii figs. 32a-6] Ordovician (Upper) U.S.S.R. (Arctic), Illaenus indeterminatus Walcott compared with G. p. 36, Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Goldillaenus Schindewolf referred to Scutelluidae (Goldillaeninae) subfam. n. p. 35, Balashova (2).

Gonioteloides Kobayashi 1955 discussed and relation to Psalikilus suggested, G. sp. cf. G. monoceros K. p. 117 pl. xxi figs. 8-9, Berg & Ross.

Goniotelus striutus n. sp. [nom. nud.] p. 77 Ordovician China, Lu.

Gourdonia Pillet 1954 discussed and referred tentatively to Asteropyginae p. 47, G. gourdoni (Barrois 1884) discussed text-fig. 1 [after Barrois], G. destombesi sp. n. p. 48 text-figs. 2-3 Devonian (Middle) France (H. Garonne), Struve Senckenbergiana leth. 40 1-2 1959.

Granularia discussed p. 101ff., G. protolenorum Lermontova text-figs. 1 (reconstr.)-2, Suvorova (1).

Granularia range in Altai-Sayan pl. i, Zhuravleva & Repina.

Granulariidae discussed and referred to Dikelocephaloidea, morphological resemblances to Asaphids may imply affinity p. 101ff., Suvorova (1).

Gravicalymene sp. nov. [undescr.] p. 208, Dean (1).

Griffithides cf. brevicaudata [sic] Gheyselink p. 207 text-fig., Yanagisawa.

Griffithides cervilatus Web., G. ? praepermicus W., G. lutugini W. ?, G. grünwaldti Moell. var. ivanovi W. pl. xii fig. 6, G. (Giphinium [recte Cyphinium]) kumpani W. var. planiloba W. listed p. 144, Ivanova.

Harpes sp. cf. macrocephalus Goldfuss [for Ferronnière 1920–22 pl. 5 fig. 99] p. 103, Pillet (2).

Harpes neogracilis R. et E. Richter 1924 p. 104 pl., Strnad (2).

Harpidae [not Harpedidae] so spelt p. 165n, Henningsmoen.

Harpidae see Harpina subord.

Harpides rugosus (Sars & Boeck) p. 274, Tjernvik.

Harpides rugosus (Sars & Boeck 1838) with cephalic doublure showing two girders p. 166 pl. ii figs. 5-11, Henningsmoen.

Harpididae Raw 1949 so spelt [not Harpidedidae], considered possibly related to Harpidae and Hungaiidae p. 164, **Henningsmoen**.

Harpididae see Harpina subord. n.

He (Bar pl. : (B. Vand of T base

[19

He

Harr

1959

Pole pole H

Bal

H

H

Horher p. 1

1

defi 9afigi pl.

H. fig Ba

pl.

p.

K

re

Harpina subord. n. [Ptychopariida] p. 415 for Harpidae, Harpididae and Entomaspididae, Whittington Treatise on Invertebrate Paleontology Part O 1959.

Heliomeroides teres p. 384, Whittington,

Hemiarges Gürich 1901 discussed p. 149, H. ambiguus (Barrande 1846) lectotype chosen and figured p. 150 pl. xi figs. 4-12, text-figs. 39-40, H. heteroclytus (B. 1846) p. 152 pl. vi figs. 7-12, text-figs. 41-43, Vanék (2).

Holdenia typa Cooper compared with young stages of Taimyraspis taimyrica gen. et sp. n. and considered based on immature specimens p. 24, Balashova (1).

Holia secristi p. 384, H. cimelia p. 385, Whittington.

Holmia cephalon pl. i fig. 4 pars, Samsonowicz (3).

Holmia kjerulfi (Lnrs. 1871) p. 447 pl. i figs. 1-10, pl. ii figs. 1-11, H. panovoi sp. n. p. 449 pl. ii figs. 12a-b, H.? sp. p. 450 pl. iii fig. 13 Cambrian (Lower) Poland (H. Cross Mts.), Samsonowicz Bull. Acad. polon. Sci. [sci. chim., etc.] 7 6 1959.

Holotrachelacea see Illaenina subord, n.

Holotrachelus punctilosus (Tornquist) 1884 p. 299 pl. lxxix figs. 3a-b, Lesnikova & Weber.

Homagnostus kasachstanicus Bal. [not seen] p. 144, Balashova & Balashov.

Homalonotus knightii König p. 477, Holland.

Homalonotus knightii König p. 514, Walmsley.

Homalonotus [misspelt Homanolotus] (Digonus) rhenanus (Koch), H. (D. ?) collini (Renaud), H. sp. p. 19, H. (D.) of. r. p. 20, **Pillet** (1).

Homalonotus sp. p. 338, Paulus.

Homalonotus (Brongniartella) sp. p. 78, Lu.

Homalops Remelé 1885 (non Motschulsky 1850) see Calyptaulax (Subgen. A).

Homolichas Schmidt specific key p. 302ff., H. deflexa (Sjögren) 1854 p. 302 pl. lxxx figs. 7a-b, 9a-b, H. depressa (Angelin) 1858 p. 303 pl. lxxx figs. 10a-12, H. eichwaldi (Nieszkowski) 1857 p. 303 pl. lxxx figs. 6a-b, 8, Lesnikova & Weber.

Homotelus taimyricus sp. n. p. 34 pl. ii figs. 15-18, H. pregradnicus sp. n. p. 36 pl. i figs. 16-17, pl. ii figs. 10, 19 Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 14 1959.

? Homotelus n. sp. [undeser.] p. 36, Thorsteinsson.

Honanaspis subgen, n. see Proasaphiscus.

Hoplolichas conicotuberculatus (Nieszkowski) 1858 p. 302 pl. lxxxii figs. 1a-2, Lesnikova & Weber.

 $Hungaia\ magnifica\ (Billings)\ from\ Maryland\ p.\ 388$  pl. liv fig. 15, Rasetti.

Hungaiidae referred tentatively to Olenina p. 158, Harpididae may be related p. 166, Dikelokephalininae a subfamily p. 161, Henningsmoen.

Hypagnostus parvifrons, H. cf. truncatus p. 12, Kindle & Whittington.

Hypermecaspididae Harrington & Leanza 1957 considered a subfamily of Olenidae, *Tropidopyge* not referred p. 161, Henningsmoen.

Hypermecaspis rugosa (Brøgger) so referred p. 174, Henningsmoen.

Hystricurus binodosus Weber 1948 p. 274 pl. lxv figs. 3a-c, Lesnikova & Weber.

Hystricurus conicus (Billings) p. 144, Balashova & Balashov.

Hystricurus oculilunatus Ross p. 112 pl. xxi fig. 2, H. ? sp. aff. H. ? genacurvus (Hintze) p. 112 pl. xxi figs. 21, 23, Berg & Ross,

Ianashtycgolia see Sanashtycgolia.

Ichangolithus ichangensis [nom. nud.] Ordovician China table 1 [p. 9], Lu.

Idahoia Walcott1925 discussed with diagnosis p. 420, I. serapio W. p. 417 pl. lx figs. 28–36, I. visconsensis (Owen) [not a geographical variant of I. s.] with immature stages p. 417 pl. lix figs. 12–32, I. latifrons (Shumard) types lost and sp. unrecognisable p. 421, Idahoia (Meeria) Frederickson 1949 so referred and discussed p. 419, I. (M.) modesta sp. n. Cambrian (Upper) U.S.A. (Idaho) p. 420 pl. lix figs. 33–45, I. (M.) undeser. sp. from Texas p. 420, Lochman & Hu J. Paleont. 33 3 1959.

Idahoia wisconsensis (Owen) p. 111 pl. xxii fig. 10, I. sp. p. 111 pl. xxii figs. 7, 11, Berg & Ross.

Idahoiidae see Olenina.

Idiomesus intermedius sp. n. p. 393 pl. li figs. 25–26 Cambrian (Upper) U.S.A. (Md & Pa), Rasetti J. Paleont. 33 3 1959.

Illaenacea see Illaenina subord, n.

Illaenidae see Ectillaeninae subfam. n.

Illaenids p. 385ff., Whittington.

Illaenina subord. n. [Ptychopariida] p. 365 for Illaenacea, Holotrachelacea and Proetacea, Jaanusson Treatise on Invertebrate Paleontology Part O 1959.

Illaenus Dalman 1827 specific key p. 291 ff., I. schmidtii Nieszkowski 1857 p. 292 pl. lxxvi figs. 4a-c, pl. lxxvii figs. 9, I. esmarkii (Schlotheim) 1826 p. 292 pl. lxxvi figs. 5a-6e, I. convexicollis Weber 1948 p. 293 pl. lxxv figs. 8-9, I. funiculatus W. 1948 p. 293 pl. lxxviii figs. 6a-b, I. oculosus Holm 1886 p. 293 pl. lxxviii figs. 3-4, I. sinuatus H. 1886 p. 293 pl. lxxviii figs. 3-4, I. sinuatus H. 1886 p. 294 pl. lxxviii figs. 3-4, I. sinuatus H. 1886 p. 294 pl. lxxviii figs. 3a-c, I. jevensis H. 1886 p. 294 pl. lxxviii figs. 2, I. talasicus W. 1948 p. 294 pl. lxxviii figs. 3a-c, I. of (Angelin 1854) p. 295 pl. lxxviii figs. 3a-c, I. of proma tukersiana H. pl. lxxviii figs. 3a-b, 4-6, I. o. forma kukersiana H. pl. lxxviii figs. 7a-8e, I. o. forma itferensis H. pl. lxxviii fig. 3, I. revaliensis H. 1886 p. 295 pl. lxxxiii figs. 1-3, I. dalmani Volborth 1863 p. 295 pl. lxxviii figs. 1-3, I. dalmani Volborth 1863 p. 296 pl. lxxvii figs. 2a-b, I. sulcifrons H. 1886 p. 296 pl. lxxvii figs. 8a-b, I. intermedius H. 1886 p. 296 pl. lxxviii figs. 4-5, I. linnarssoni H. 1883 p. 296 pl. lxxviii figs. 4-5, I. linnarssoni H. 1883 p. 296 pl. lxxviii figs. 1a-c, I. oviformis Warburg 1925 p. 297 pl. lxxviii figs. 6a-e, Lesnikova & Weber.

Illaenus aff. aemulus Salter (=I. namhsinensis Reed) p. 1548, Bordet et al.

Illaenus americanus Billings, I. groenlandicus Troedsson, I. sp. p. 36, Thorsteinsson.

[19

KL

Kl

Ke

Ke

Carn

Repi

K

K. n

sp. 1

p. 2 p. 2

pl.

Ress

& R

She

pale

Zh

for

Car

et a

Ka

no

be

Ca

pl.

 $\mathbf{z}$ 

ELL

th

(I

Can

F

I

K

Zhu

of Sa

Illaenus aff. americanus Billings p. 88, I. ef. angusticollis B. p. 91, Colquhoun.

Illaenus angustifrons Holm p. 6ff., I. mascei H., I. roemeri Volb. p. 11, Männil.

Illaenus huangi p. 73, I. deltoides p. 77 n. spp. [nom. nuda] Ordovician China, Lu.

Illaenus cf. thomsoni Salter p. 361, Miroshnikov et al.

Illaenus valvulus Raymond pl. iii figs. 5-6, I. latiaxiatus Bradley et R. pl. iii figs. 7-9, I. depressicapitatus B. pl. iii figs. 10-14, I. sp. pl. ii fig. 13, Balashoya (1).

Illaenus valvulus Raymond 1925 p. 38, I. latiaxiatus Raymond et Narraway p. 40, I. depressicapitatus Bradley 1930 p. 39, I. sp. p. 41; I. indeterminatus Walcott referred to Goldillaeninae subfam. n. p. 35 and compared with Goldillaenoides gen. n. p. 36, Balashova (2).

Inouyella Resser & Endo referred to Namanoiidae p. 222, Chang.

Inouyia fongfongensis sp. n. [nom. nud. 1957 when referred to Inoyellaspis] p. 211 (236) pl. iv fig. 19 Cambrian (Middle) China, Chang Acta palaeont. sinica 7 3 1959.

Inouyina Poletaeva 1936 referred to Namanoiidae but probably a synonym of Taitzuia p. 223, Chang.

Inouyina Poletaeva discussed and referred to Namanoidae p. 141, I. quadratica P. lectotype selected [illegitimate] p. 143 pl. ix figs. 1-6, Pokrovskaya.

Inouyina [misspelt Snouying on pl.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Inoyellaspis fongfongensis Chang 1957 nom. nud. see Inouyia fongfongensis sp. n.

Iranaspidinae subfam. n. [Leiostegiidae] p. 315 for Iranaspis and Plethopeltella, Lochman[-Balk] Treatise on Invertebrate Paleontology Part O 1959.

Iranaspis see Iranaspidinae subfam, n.

Irvingella text-fig. 3e, Kindle & Whittington.

Irvingella norilica Las. [not seen] p. 359, Miroshnikov et al.

Isalaux stricta Kram. [not seen] p. 360, Miroshnikov et al.

Isocolidae—diagnosis discussed and Thomondia referred p. 377, Weir.

Isocolidae see Taimyraspis gen. n.

Isocolus aff. sjögreni Angelin 1854 p. 299 pl. lxxix figs. 4a-b, Lesnikova & Weber.

Isoteloides homalonotoides (Walcott) var. taimyrica var. n. p. 33 pl. ii fig. 4 Ordovician (Upper p. 33 but Middle pl. expl.), I. kimmswickensis Bradley var. nordica var. n. p. 34 pl. ii fig. 14 Ordovician (Upper) U.S.S.R. (Arctic), Balashova Sborn. stat. paleont. biostrat. 14 1959.

Isotelus p. 89n, Thorsteinsson.

Isotelus aff. iowaensis (Owen), I. cf. latus Raymond p. 85, I. gigas DeKay p. 91, Colquhoun.

Isotelus aff. iowensis Owen 1852 p. 31 pl. ii figs. 7-9, I. sp. 1 p. 32 pl. ii fig. 15, I. sp. 2 p. 33 pl. ii fig. 3 Ordovician (Upper) U.S.S.R. (Arctic), Balashova (1).

Isotelus platyrhachis (Steinh.) p. 6ff., Männil,

Isotelus remigium (Eichwald) 1860 p. 281 pl. lxxiii fig. 5, I. platymarginatus Raymond 1910 p. 281 pl. lxxiii fig. 3, I. aktchokensis Weber 1948 p. 282 pl. lxxiii fig. 2, Lesnikova & Weber.

Isotelus sp. p. 387, Whittington.

Jakutidae fam. n. [Corynexochoidea] for Jakutus, Bathyuriscellus, Judaiella, Parapoliella, Malykania and Lenaspis gen. n. p. 66, Suvorova Paleont. Zhurnal 1959 3.

Jakutus see Jakutidae fam. n.

Jakutus quadriceps Lerm. p. 1372, Z. A. Zhuravleva (2).

Johoaspis [nom. nud.] sp. p. 5, Lazarenko.

Judaiella see Jakutidae fam. n.

Judomia see Callaviinae subfam. n.

Kadyella gen. n. [Protoleninae] p. 58 for type K. ubsanurica sp. n. p. 59 pl. i figs. 8-9, 13-16, pl. ii figs. 1-2, K. kadyensis sp. n. p. 61 pl. i fig. 10 Cambriau (Lower) U.S.S.R. (Tuva & Yakutia), K. sp. p. 32, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Kainella alimbetica Bal. [not seen] p. 144, Balashova & Balashov.

Kainella sp. cf. K. billingsi Walcott p. 117 pl. xxi fig. 26, pl. xxii figs. 6, 13, Berg & Ross.

Kainella lohanpoensis Sheng n. sp. [nom. nud.] p. 36 Ordovician (Lower) China, Lu.

Kameschkoviella Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Karakolia gen. et sp. nov. (Chernysheva) [nom. nud.] p. 81, Zhuravleva et al. (1).

Kazelia norilica Las. [not seen] p. 359, Miroshnikov et al.

Keithia intermedia sp. n. p. 392 pl. liv figs. 23-27, K. sp. undet. p. 392 pl. li fig. 24 Cambrian (Upper) U.S.A. (Maryland), Rasetti J. Paleont. 33 3 1959.

Keithiella text-fig. 3a, Kindle & Whittington.

Keithiella cylindrica (Billings) p. 391 pl. li fig. 23, K. sp. undet. p. 392 pl. lii figs. 17–18, **Rasetti.** 

Kendallia Berg 1953 non Evermann & Shaw 1927 see Kendallina nom. n.

Kendallina gen, n. [Parabolinoididae] (nom. n. for Kendallia Berg 1953 non Evermann & Shaw 1927) p. 272, R. R. Berg Treatise on Invertebrate Paleontology Part O 1959.

Kijanella range in Altai-Sayan pl. i, Zhuravleva & Repina.

Kingstonia sp. p. 359, Miroshnikov et al.

Kjerulfia see Callaviinae subfam. n.

Kjerulfia  $\ell$  lagowiensis Czarnocki [sp. nondeser.] p. 450 pl. iii figs. 10-12, K. sp. p. 450 pl. iii figs. 1-9, Samsonowicz (1).

ii

2

1

i

1

n

Klotziella poletaevae [poletayevi] Lermontova type of Sayanella gen. n. [quod vide] p. 150, Pokrovskaya.

Kloučekia solitaria (Barr.) p. 48, Vaněk (3).

Kochina see Trachycheilus gen. n.

Kochina ? angustata sp. n. p. 212 pl. xxiii fig. 3 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg, Instn No. 563 1945.

Kolbinella prima Rep. [nom. nud.], K. sp. p. 77, Zhuravleva et al. (1),

Kootenia range in Altai-Sayan pl. i, Zhuravleva & Repina.

Kootenia tetraspinosa sp. n. p. 199 pl. xx figs. 1-4, K. mckeei sp. n. p. 200 pl. xx figs. 5-10, K. schenki sp. n. p. 200 pl. xx figs. 11-17, K. simplez sp. n. p. 201 pl. xxii figs. 16pars-17, K. havasuensis sp. n. p. 202 pl. xxii figs. 14-17, K. sp. undet. p. 202 pl. xxii figs. 14-17, K. sp. undet. p. 202 pl. xxii figs. 16pars-17, Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Kootenia sp. p. 1033, Zhuravleva et al. (2).

Kooteniella range in Altai-Sayan pl. i, Zhuravleva & Repina.

Kooteniella slatkowskii Smidt p. 626, Anatol'eva & Shelkovnikov.

Koptura longibiloba sp. n. p. 209 (232) pl. iv fig. 2 Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Kounamkites sp. p. 5, Lazarenko.

Krolina pressulata Rep. [nom. nud.] p. 71, Zhuravleva et al. (1).

Kujandaspis gen. n. [no family assigned] p. 110 for type K. kujandensis sp. n. p. 111 pl. i fig. 5 Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Kujandaspis kujandensis Ivshin p. 359, Miroshnikov et al.

Kujandina gen. n. [no family assigned] p. 111 for type K. taskudukensis sp. n. p. 112 pl. i fig. 8 Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Kuraspis obscurus gen. et sp. n. (N. E. Chernysheva) [nom. nud.] p. 651, Petrakov.

Labradoria [angustifrons] Ivshin 1957 does not belong to L. p. 131n, Pokrovskaya.

Ladadiscus gen. n. [Eodiscinae] p. 167 for type L. limbatus sp. n. p. 168 pl. xi figs. 5-8, 10-15, 17 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Laminurus Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Laminurus inornata Repina [nom. nud.] p. 1032, Zhuravleva et al. (2).

Lapidaria see Niobinae subfam. n.

Laticephalus gen. n. [Incertae sedis, compared with Edelsteinaspis and Paleofossus] p. 158 for type L. trapezoidalis sp. n. p. 159 pl. viii figs. 10, 13, L. tuberosus sp. n. pp. 159, 161 pl. x figs. 1-2 [but these figs. referred to trapezoidalis p. 159] Cambrian (Lower) U.S.S.R. (Tuva etc.), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Laticephalus range in Altai-Sayan pl. i, Zhuravleva & Repina.

Lauzonella see Loganellidae fam. n.

Leiagnostus alimbeticus Bal. [not seen] p. 144, Balashova & Balashov.

Leimitzia Sdzuy 1955 discussed and referred to Dikelokephalininae p. 162, **Henningsmoen**,

Leiocoryphe cf. L. gemma Clark p. 385 pl. liii fig. 32, L. transversa Rasetti [including L. n. sp. of R. 1945] with median suture on cephalic doublure p. 385 pl. liii figs. 25-30, L. cf. L. occipitalis R. p. 386 pl. liii fig. 31, L. cf. L. brevis p. 378, Rasetti.

Leiostegiidae see Iranaspidinae subfam. n.

Leiostegium Raymond 1913 discussed p. 113, L. (Leiostegium) manitouensis Walcott p. 114 pl. xxi figs. 10-13, 17, 24, L. (L.) sp. p. 114 pl. xxii figs. 10-13, 17, 24, L. (L.) sp. p. 114 pl. xxii figs. 1-2, L. (Perischodory) R. 1937 so considered p. 114, L. (P.) incompertum sp. n. p. 115 pl. xxii fig. 15, pl. xxii fig. 4, L. undet. cranidia [may belong to L. (L.) m. or L. (P.) i.] p. 115 pl. xxii figs. 14, 19, L. (Manitouella) [Ulrich MS] subgen. n. for type L. (M.) ulrichi sp. n. p. 115 pl. xxi figs. 1, 3, 4, 6, pl. xxii figs. 5, 9 Ordovician (Lower) U.S.A. (Colorado), Berg & Ross J. Paleont. 33 1 1959.

Leiostegium ? sp. undet. from Cambrian (Upper) p. 413 pl. lvii figs. 21-22, Lochman & Hu.

Lenaspis gen. n. [Jakutidae] for type L. opipara sp. n. p. 68 pl. vi figs. 1-3 text-figs. 1a-6, L. limata sp. n. p. 69 pl. vi fig. 4 text-figs. 2a-6 Cambrian (Lower) U.S.S.R. (Yakutia), Suvorova Paleont. Zhurnal 1959 3.

Lermontovia Suvorova 1956 a synonym of Protolenus to which L. lenaica and L. convexa S, are referred p. 42, Pokrovskaya.

Lermontoviella gen. n. [Antagmidae] p. 154 for the L. shanganica sp. n. p. 155 pl. ix figs. 7-9, L. sayanica sp. n. pp. 155, 157 for Psychoparia? sp. Poletaeva 1936 Cambrian (Lower) U.S.S.R. (Tuva & Sayan), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Lermontoviella range in Altai-Sayan pl. i, Zhurav-leva & Repina.

Lermontoviidae Suvorova 1956 discussed and so considered, taken to include *Lermontovia*, *Bergeroniellus*, *Bergeroniaspis* and *Olekmaspis* p. 395,

Lermontoviinae Suvorova a synonym of Protoleninae [quod vide] p. 39, Pokrovskaya.

Lermontoviinae see Lermontoviidae.

Levisella see Loganellidae fam. n.

Levisella nasuta sp. n. p. 394 pl. liv figs. 1–3 Cambrian (Upper) U.S.A. (Maryland), Rasetti J. Paleont. 33 3 1959.

Levisia Walcott 1911 referred to Namanoiidae p. 222, L. adrastia (W.) so referred and discussed p. 225 text-figs. 18-19, Chang.

Liaoyangaspis gen. n. [nondescr. 1957] [no family ascribed] p. 197 (219) for type Asaphiscus bassleri Resser & Endo 1937 (stated to have 11 thoracic segments) with synonym A. kobayashii R. & E. [given as

type of L. in 1957] p. 198 (220) pl. ii figs. 1-3 text-figs. 5-6, L. walcotti (R. & E.) restr. text-fig. 7, L. tsutsumii (E.) text-fig. 10, L. endoi sp. n. for A. walcotti R. & E. 1937 partim (pl. 43 fig. 6) text-fig. 8, L. wangi sp. n. for A. walcotti of Lu 1957 text-fig. 9 p. 200 (221) Cambrian (Middle) China & Manchuris, L. peiensis (R. & E.) also referred p. 220, Chang Acta palaeont. sinica 7 3 1959.

Lichaidae [so spelt] Hawle et Corda 1847 morphology and phylogeny discussed p. 82ff., discussed and classified into Lichainae and Ceratarginae p. 89, Vaněk (2).

Lichainae [so spelt] Hawle et Corda 1847 discussed and taken to include Platylichas, Dicranogmus, Dicranopeltis, Lichas, Amphilichas, Conolichas, Hoplolichas, Leiolichas, Uralichas, Arctinurus, Gaspelichas, Terataspis and Ceratolichas p. 91, Vaněk (2).

#### Lichakephalidae see Lichida ord. n.

Lichas Dalman 1827 discussed with synonym i. a. Metopolichas p. 92, L. praecursor Holub 1911 holotype refigured p. 94 pl. i fig. 8 text-fig. 1, L. incola Barrande 1856 lectotype selected and figured including L. uvus B. 1872 partim (pl. 5 fig. 23, pl. 6 figs. 25-26, pl. 10 fig. 17—see also Uralichas) p. 96 pl. iii figs. 1-7 text-figs. 2-5; L. sp. Růžička 1926 pl. 3 figs. 4-5 referred to Platylichas kloučeki p. 115, L. parvula Novák 1890 a synonym of Acanthopyge parva parva p. 125, Vaněk (2).

Lichas sp. p. 339, L. (Ceratarges) sp. Gruppe armatus (Goldfuss) p. 353, Paulus.

Lichida ord. n. p. 495 for Lichidae and Lichakephalidae, Moore Treatise on Invertebrate Paleontology Part O 1959.

Lichidae see Lichida ord. n.

Lichidae see Lichaidae.

Lichids p. 385ff., Whittington.

Lichinae see Lichainae.

Lioharpes venulosus Corda p. 103, Pillet (2).

Litagnostus sp. undet. p. 381 pl. li fig. 7, Rasetti.

Lobopyge Přibyl et Erben 1952 discussed p. 127, L. branikensis (Barrande 1872) lectotype refigured p. 128 pl. vii figs. 9-11, text-figs. 18-21, L. pragensis (Bouček 1933) holotype refigured p. 132 pl. vii figs. 8, pl. viii figs. 8-9, text-figs. 22-23, L. richteri sp. n. p. 133 (157) pl. viii figs. 4-7, text-figs. 24-26 Silurian (Upper), L. docekali [dočekali on pl. expl.] sp. n. [including Acanthopyge aff. haueri of Chlupáč & Vaněk 1957 p. 125] p. 136 (158) pl. viii figs. 1-3 text-figs. 27-29 Devonian (Lower) Czechoslovakia, Vaněk Bohemia Centralis [A] 1 3 1959.

Loganellidae fam. n. [Remopleuridacea] p. 331 for Loganellus, Lauzonella and Levisella, Rasetti Treatise on Invertebrate Paleontology Part O 1959.

Loganellus-group see Olenina.

Loganellus see Loganellidae fam. n.

Loganellus cf. L. similis Rasetti p. 394 pl. li figs. 28-29, Rasetti.

Loganopeltis nannus Bal. [not seen] p. 144, Balashova & Balashov.

Loganopeltoides text-fig. 3d, Kindle & Whittington.

Lonchodomas carinatus Cooper 1953 morphology and ontogeny p. 473 pl. xxxii figs. 1-23, pl. xxxiii figs. 1-32 text-figs. 8A-c (reconstrs.), Whittington.

Lonchodomas pennatus (La Touche), L. aff. rostratus (Sars) p. 207, L. cf. p. p. 208, Dean (1).

Lonchodomas rostratus (Sars et Boeck) var. taimyrica var. nov. pl. vi figs. 6-7, Balashovs (1).

Loshanella loshanensis n. gen. n. sp. [nom. nud.] p. 39, L. ? sp. p. 37 Ordovician (Lower) China, Lu.

Lotagnostus cf. L. trisectus (Salter) p. 381 pl. li figs. 8-9, Rasetti.

Luia gen. n. [related to Anomocarellidae] p. 193 (215) for type L. typica sp. n. [nom. nud. 1957] p. 194 (216) pl. i figs. 1-4 text-fig. 1, L. yaochiayuensis sp. n. p. 194 (216) pl. i figs. 5-7, L. shantungensis sp. n. p. 194 (217) pl. i fig. 8 Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Lunolenus n. g. (nom. nud.) [Doleroleninae] p. 396, L. lunae n. sp. (nom. nud.) pl. i fig. 2 (reconstr.), Sdruy.

Lyralichas bronnikovi Weber 1932 p. 304 pl. lxxx figs. 15a-c, Lesnikova & Weber.

Malykania see Jakutidae fam. n.

Malykania gribovae Suv. p. 797, Z. A. Zhuravleva (1).

Malykania gribovae Suv. p. 1372, Z. A. Zhuravleva (2).

Manchuriella Resser & Endo discussed and considered probably a subgenus of Proasuphiscus p. 227; M. pustulosa R. & E. restr. referred to Szeaspis gen. n. p. 230, Chang.

Manitouella subgen, n, see Leiostegium.

Matthewlenus Hupé a synonym of Protolenus p. 42, Pokrovskaya,

Meeria Frederickson 1949 discussed and considered a subgenus of Idahoia [quod vide] p. 419, Lochman & Hu.

Meeria livae sp. n. error pro Meeria lirae sp. n. Frederickson 1949, Z. R. 86 11 (for 1949) 1951.

Megalaspis Angelin specific key p. 278ff., M. limbata (Boeck) 1837 p. 279 pl. lxvi figs. 3–4, M. gibba Schmidt 1898 p. 279 pl. lxvii figs. 1a–b, M. hyorhina (Leuchtenberg) 1843 p. 279 pl. lxvii figs. 6a–c, M. h. var. kolenkoi S. p. 279 pl. lxvii fig. 5, M. planilimbata A. 1852 p. 279 pl. lxvii figs. 4–5, M. acuticauda A. 1854 p. 280 pl. lxv fig. 11, M. a. var. triangularis S. 1904 p. 280 pl. lxvii figs. 3a–b, M. centaurus var. rudis A. 1854 p. 281 pl. lxvii fig. 2, Lesnikova & Weber.

Megalaspis limbata (Boeck 1837) from glacial boulder p. 40 text-fig., Strnad (5).

Megalaspis limbata (Boeck 1837) p. 322 pl. xxv figs. 1-2, pl. xxvi fig. 3, Strnad (8).

Megalaspis obtusicauda Bohlin, M. aff. gigas Angelin (=M. centaurus Schmidt) recorded p. 138, Balashova & Balashov.

Mendospidella quebradensis Rusc. discussed p. 6, Rusconi. Mer pis g S.S.S.

[195

Mer

type i (Lowe Trud.

Me Remo

Me

Me referre

diseu pl. i Me Dole

with sp. n pl. i i for M. i belo Char

M. d M. 2a-l figs. pl. 1 figs. pl. 1

p. 9

1

M. info

for p. Pr

A K

Menevia of. venulosa text-fig. 3i, Kindle & Whittington.

Menneraspidae fam. n. [Redlichoidea] for Menneraspis gen. n. p. 83, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Menneraspis gen. n. [Menneraspidae] p. 84 for type M. striatus sp. n. p. 85 pl. iii figs. 1–10 Cambrian (Lower) U.S.S.R. (Tuva & Yakutia), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Menoparia—ontogeny compared with that of Remopleurides p. 397ff., Whittington.

Mesotaphraspis inornata, M. parva p. 384 ff., Whittington.

Metadoxides Bornemann 1891 discussed and referred to Doleroleninee, Anadoxides Matthew considered a synonym, M. torosus pygidial structure discussed, M. richterorum n. sp. (nom. nud.) p. 397 pl. i figs. 3-4 (reconstrs.), Sdzuy.

Metadoxidinae Whitehouse 1939 a synonym of Doleroleninae p. 396, **Sdzuy**.

Metanomocarella gen. n. [nondeser. 1957] [compared with Anomocarella and Aojia] for type M. rectangula sp. n. Cambrian (Middle) China (Shantung) p. 195 (217) pl. i figs. 9-12 text-fig. 2, M. tumida (Endo) apparently for Anomocarella tumida E. 1944 [stated to be type of M. in 1957] to which (or to another sp. of M.) probably belongs Aojia tumida [Resser & Endo 1937 p. 218, Chang Acta palaeont. sinica 7 3 1959.

Metopolichas Gürich specific key p. 300ff., M. celorhin (Angelin) 1854 p. 301 pl. lxxxi figs. 5a-c, M. verrucosa (Eichwald) 1843 p. 301 pl. lxxx figs. 2a-b, M. kuckersiana (Schmidt) 1885 p. 301 pl. lxxx figs. 5a-b, M. pachyrhina (Dalman) 1828 p. 301 pl. lxxxi figs. 6a-7, M. p. var. longerostrata S. pl. lxxxi figs. 8a-b, M. anderkenensis [sic] Weber 1948 p. 302 pl. lxxx figs. 1a-b, Lesnikova & Weber.

Metopolichas Gürich 1901 a synonym of Lichas p. 92, Vaněk (2).

Micmacca sp. p. 30, Pokrovskaya.

Microdiscus parkeri Walcott p. 124 (pl. 50) fig.,

Minkella gen. n. [Dokimocephalidae] for type M. americana sp. n. with immature stages and inferred sexual dimorphism p. 414 pl. lviii figs. 1-3, 5, 7-15, 18-20 [2 forms with occipital spine, including holotype], p. 416 pl. lviii figs. 4, 6, 16-17 [3 forms without o. spine] Cambrian (Upper) U.S.A. (Idaho); Taenicephalus megalops Kobayashi some cranida may belong to Minkella p. 416, Lochman & Hu J. Paleont. 33 3 1959.

Mitchellaspis gen. n. [? Encrinurinae] (nom. n. for Mitchellia Vogdes 1917 non de Koninck 1877) p. 447, Henningsmoen Treatise on Invertebrate Paleontology Part O 1959.

Mitchellia Vogdes 1917 non de Koninck 1877 see Mitchellaspis nom. n.

Monorakos lopatini Schmidt 1886 p. 312 pl. lxxxiv fig. 12, Lesnikova & Weber.

Monorakos złobini sp. nov. pl. iv figs. 1-2, M. nordicus sp. nov. pl. iv figs. 3-4, M. magnus Kramarenko pl. iv fig. 5, M. mutabilis K. pl. iv fig. 6, Balashova (1). Mrassina submissa Repina [nom. nud.] p. 1032, Zhuravleva et al. (2),

Mundocephalina range in Altai-Sayan pl. i, Zhuravleva & Repina.

Namanoia ? grandis sp. nov. [nom. nud.] p. 29, Pokrovskaya.

Namanoia namanensis Lerm. p. 798, Z. A. Zhuravleva (1).

Namanoidae—Inouyina referred p. 141, Pokrov-skaya.

Namanoiidae Lermontova taken to include Namanoia, Ordosia, Taitzuia, Inouyella, Inouyina, Levisia, and Poshania & Peichiashania genn. n. p. 222, Chang.

Neoaldanaspis aff. arcticus Las. [not seen] p. 359, Mikoshnikov et al.

Neosynphoria see Synphoria.

Nieszkowskia cephaloceras (Nieszkowski) 1857 p. 310 pl. lxxxiii figs. 17-19, Lesnikova & Weber.

Nileus armadillo Dalman 1827 p. 290 pl. lxxii fig. 13, N. tengriensis Weber 1948 p. 290 pl. lxxii figs. 14a-b, Lesnikova & Weber.

Nileus liangshanensis n. sp. [nom. nud.] p. 76 Ordovician China, Lu.

Nileus limbatus Brögger p. 144, Balashova & Balashov.

Nileus limbatus Brøgger listed p. 174, Henningsmoen.

Ninaspis gen. n. [no family assigned] p. 113 for type N. tchernyshevae gp. n. p. 114 pl. i fig. 4 Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshim Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Niobe see Niobinae subfam. n.

Niobe insignis Linnarsson listed p. 174, Henningsmoen.

Niobe insignis Linnarsson p. 274, Tjernvik.

Niobe insignis Linn., N. laeviceps Dalm. p. 144, Balashova & Balashov.

Niobe laeviceps (Dalman) 1827 p. 281 pl. lxxii figs. 11–12, N. frontalis (D.) 1827 p. 281 pl. lxxii fig. 10, Lesnikova & Weber.

Niobella see Niobinae subfam. n.

Niobella obsoleta (Linnarsson) listed p. 174, Henningsmoen.

Niobina see Niobinae subfam. n.

Niobinae subfam. n. [Asaphidae] p. 350 for Niobe, Bohemopyge, Lapidaria, Niobella, Niobina, Norinia and Yuepingia, Jaanusson Treatise on Invertebrate Paleontology Part O 1959.

Nodiceps gen. n. [Edelsteinaspididae] p. 71 for type N. onerosa sp. n. p. 72 pl. vi figs. 5–9 text-figs. 3a-2, 4 Cambrian (Lower) U.S.S.R. (Yakutia), Suvorova Paleont. Zhurnal 1959 3.

Norinia see Niobinae subfam. n.

Norwoodiacea see Bolaspididae fam. n.

Novaspis aff. albida (Reed) including Trinucleus concentricus of Baily 1862 p. 371 pl. lxii figs. 5-8,

Odontochile Hawle & Corda 1847 placed on Official List of Generic Names, gender deemed feminine, designation of Asaphus hausmanni Brongniart 1822 [quod vide] as type p. 43, 0. applanatus H. & C. 1847 [considered nom. dub. by R. P. Tripp & W. F. Whittard p. 53] Vogdes' 1925 selection as type of O. set aside, Opinion 537.

Odontochile spinifera Barrande p. 22 text-fig. 3, Suñer Coma & Fernández de Villalta.

Odontopleurida ord. n. p. 504 for Odontopleuridae and Eoacidaspididae, Whittington Treatise on Invertebrate Paleontology Part O 1959.

Odontopleuridae see Odontopleurida ord. n.

Ogygia sola Barrande referred to Bronteopsis p. 215, Vaněk (1).

Ogygia sp. v. Freyberg 1923 may belong to Phacopacea p. 41, Struve (1).

Ogygites primulus (Barr.) for Proetus ? p. B. p. 46, O. nobilis (B.) p. 47ff., Vaněk (3).

Olekmaspis Suvorova 1956 referred to Lermontoviidae p. 395, **Sdzuy**.

Olekmaspis Suvorova 1956 a synonym of Bergeroniaspis p. 45, Pokrovskaya.

Olenaspella sp. p. 360, Miroshnikov et al.

Olenellacea—morphology and evolution discussed p. 151ff., Stubblefield (3),

Olenellidae see Callaviinae subfam. n.

Olenellus cf. fremonti Walcott p. 193 pl. xix figs. 21–22, O. sp. [spp.] undet. p. 193 pl. xix figs. 23, 24, 25–26, Resser.

Olenellus ? problematicus sp. n. p. 157 pl. i fig. 12 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Olenidae referred to Olenina and relations to Parabolinoididae and Ptercoephaliidae [quae vide] discussed p. 157; Hypermecaspididae considered a subfam. of Olenidae p. 161, Henningsmoen.

Olenina Swinnerton 1915 so accredited and taken to include Olenidae, Pterocephaliidae, Parabolinoididae, Idahoiidae, Elviniidae, Irvingella, Dunderbergia, Conaspis, Comanchia, Ptychaspididae, Saukiidae, Dikelocephalidae, ? Hungaiidae, ? Loganellus-group, ? Remopleurididae, ? Andrarinidae, ? Anomocaridae p. 158, Henningsmoen.

Olenoides ex gr. convexus Lerm., O. cf. hibridus Resser p. 626, Anatol'eva & Shelkovnikov.

Olenoides manchuriensis Endo 1944 tentatively referred to Prodamesella gen. n. p. 219, Chang.

Olenoides nevadensis (Meek) p. 84, Bright.

Olentella gen. n. [no family assigned] p. 115 for type O. olentensis sp. n. p. 117 pl. i fig. 7 Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Olenus? Baily 1862 partim referred to Corrugatagnostus convergens sp. n. p. 370, Weir.

Onchocephalus range in Altai-Sayan pl. i, Zhuravleva & Repina.

Onchocephalus sp. p. 1032, Zhuravleva et al. (2).

Onchometopus volborthi Schmidt 1898 p. 281 pl. lxxii figs. 6-7, Lesnikova & Weber.

Onchonotus richardsoni (Walcott) p. 394 pl. li figs. 5-6, Rasetti.

Onnia gracilis (Bancroft), O. cf. superba (B.) p. 207,

Onnia ornata (Sternberg) p. 47, O. goldfussi (Barr.) p. 48, Vaněk (3).

Opsidiscidae Hupé discussed p. 177 [see also Tannudiscus gen. n.], Pokrovskaya.

Ordosia Lu 1954 referred to Namanoiidae p. 222, O. fimbricauda L. text-fig. 12, Chang.

Ormathops oriens (Barr.) p. 46, O. mirus Šnajdr p. 47, Vaněk (3).

Orometopidae see Pagometopus gen. n.

Orometopus elatifrons (Angelin) p. 274, Tjernvik.

Orometopus elatifrons (Ang.) p. 144, Balashova & Balashov.

Orometopus primigenius Størmer, O. elatifrons (Angelin) listed p. 174, Henningsmoen.

Orphanaspis sp. p. 103, Pillet (2).

Oryctocara ovata Tehern, p. 5, Lazarenko.

Oryctocephalops frischenfeldi Lerm. p. 5, Lazarenko.

Oryctocephalus (Viñakainella) asperoensis Rusc. discussed p. 6, Rusconi.

Oryctocephalus reynoldsiformis Lerm. p. 5, Laz-arenko.

Otarion see Cyphaspis.

Otarion (O.) convexum (Hawle a Corda 1847) p. 7 text-fig., Strnad (1).

Otarion (O.) ferronnierense Pillet, O. (O.) convexum Corda, O. (O.) hydrocephalum barrandei C. [respectively for Ferronniere 1920-22 pl. 1 figs. 6, 7, 8] p. 102, O. (Coignouina) nov. sp. [undescr.] near davidsoni Barrande p. 103, Pillet (2).

Otarion sp. pp. 214, 218, Dean (1).

Otarion sp. p. 19ff., Pillet (1).

Otarion sp. p. 249, Shackleton.

Otarion sp., "O." sp. p. 387, Whittington.

Otarionids p. 384, Whittington.

Ouijjania Hupé 1952 [sic] referred to Bigotininae p. 401, Sdzuy.

Ovalocephalus gen. n. [Deiphoninae] for type O. kelleri sp. n. including Staurocephalus sp. Weber 1948 p. 1316 pl. figs. 3a, 6, e, text-fig. 4e Ordovician (Middle) U.S.S.R. (Kazakhstan), Koroleva C. R. Acad. Sci. U.R.S.S. (ns) 124 6 1959.

Pachyaspis moorei sp. n. p. 210 pl. xxiii figs. 7-9, P. fonticola sp. n. p. 211 pl. xxiv figs. 31-32, P. sp. undet. p. 211 pl. xxiv figs. 16-21 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Hystri Pae Pag

[195

Pac

Pag pariso figs. I figs. I Camb

> Pag Pag Repir

Pa P. g (Tree Tidss

> P. ze P. ze 15 C Trud

& R

& Re

P

P Bala P deri

rec P.

> figs P. p. figs 7.

> > Po

80

I

pl.

P. Be p. Pachycranium ? sp. Ross 1951 may belong to Hystricurus p. 113, Berg & Ross.

Paedeumias sp. p. 1686, Theokritoff.

Pagetia ferox Lerm. p. 5, Lazarenko.

Pagetides Rasetti 1945 discussed p. 5 with comparison of spp. table 1, P. sibiricus sp. n. p. 7 pl. i figs. 1-11, P. sibiricus var. calvus var. n. p. 11 pl. i figs. 12-19, P. spinosus sp. n. p. 12 pl. i figs. 20-31 Cambrian (Middle) U.S.S.R. (N. Siberia), Lazarenko Sborn. stat. paleont. biostrat. 14 1959.

Pagetiidae see Shivelicus gen. n.

Pagetina range in Altai-Sayan pl. i, Zhuravleva & Repina.

Pagetina persubtilis Repina [nom. nud.] p. 1032, Zhuravleva et al. (2).

Pagometopus gen. n. [Orometopidae] p. 169 for type P. gibbus sp. n. p. 170 pl. ii figs. 1-4 Ordovician (Tremadoc) Norway, Henningsmoen Norsk geol. Tidsskr. 39 2-3 1959.

Palaeolenus range in Altai-Sayan pl. i, Zhuravleva & Repina.

Paleofossus gen. n. [Edelsteinaspidae] for type P. zaicevi sp. n. p. 131 pl. viii figs. 1-2, 4-5, 8-9, 12, 15 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Paleofossus range in Altai-Sayan pl. i, Zhuravleva & Replna.

Paleofossus see Laticephalus gen. n.

Palpebralia see Parapalpebralia.

Parabolina sp. nov. [undeser.] p. 222, Shackleton.

Parabolinella caesa Lake & 'P.' longispina (Belt) horizon p. 56, Stubblefield (1).

Parabolinella limitis Brögger p. 144, Balashova & Balashov.

Parabolinoididae referred to Olenina but not derived from Olenidae p. 157, Henningsmoen.

Parabolinoididae see Kendallina nom. n.

Paraceraurus aculeatus (Eichw.) text-fig. 6 [reconstr.], Kaljo (2).

Paracybeloides aff. loveni (Linnarsson) p. 207ff., P. sp. ind. p. 214, Dean (1).

Paradoxidacea see Redlichiina subord. n.

Paradoxides oelandicus Sjögren 1872 p. 441 pl. i figs. 1-4, ? 5, P. pinus Holm p. 442 pl. i figs. 6, ? 7-8, P. torelli H. p. 443 pl. ii fig. 1, P. insularis Wgård p. 443 pl. ii figs. 2-3, P. czarnockii sp. n. p. 443 pl. ii figs. 4a-b, 5, P. kozlowskii sp. n. p. 444 pl. ii figs. 6a-b, 7, P. samsonowiczi sp. n. p. 445 pl. ii figs. 8a-b, P. sp. No. 1 p. 445 pl. ii figs. 8a-b, diddle) Poland (H. Cross Mts.), Orlowski Bull. Acad. polon. Sci. [sci. chim., etc.] 7 6 1959.

Paradoxides of the ölandicus Sjögr. group p. 771 pl. figs. 1-6, P. tessini Brng. p. 771, Orlowski (1).

Paradoxides pradoanus Barr. & Vern. p. 119 (pl. 45) fig., P. spinosus Boeck p. 120 (pl. 46) fig., P. rotundatus Barr. p. 121 (pl. 47) fig., P. bohemicus Boeck p. 122 (pl. 48) 2 figs., P. rugulosus Corda p. 123 (pl. 49) fig., P. afi. r. p. 123, Badillo.

Paradoxides sp. p. 5, Lazarenko.

Paraerbia range in Altai-Sayan pl. i, Zhuravleva & Repina.

Paralejulus 800 Scutellum.

Parapalpebralia Erben gen. nov. [nom. nud.] including Armorican forms referred by Pillet 1953 to Piriproctus and Palpebralia p. 103, P. f sp. [for Ferronnière 1920–22 pl. 1 fig. 13] p. 102, Pillet (2).

Parapilekia speciosa (Dalman) listed p. 174, Henningsmoen.

Paraplethopeltis? genacurvus and P.? genarectus Hintze spp. may belong to Hystricurus p. 112, Berg & Ross.

Parapoliella see Jakutidae fam. n.

Parasolenopleura linnarssoni (Brögg.) p. 771, Orlowski (1).

Parehmania nitida sp. n. p. 207 pl. xxiv figs. 7-15, P. tontoensis sp. n. p. 207 pl. xxiv fig. 22, P. kwaguntensis sp. n. p. 208 pl. xxv figs. 9-13 partes, pl. xxv figs. 7-10, P. sp. undet. p. 209 pl. xxv figs. 3-4 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instu No. 563 1945.

Pedinocephalus gen. n. [no family assigned] p. 114 for type P. bublichenkoi sp. n. p. 115 pl. i fig. 3 Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Peichiashania Chang [nondescr. 1957] gen. n. [Namanoiidae] for type Eymekops rectangularis Resser & Endo [recte Endo] 1937 Cambrian (Upper) Manchuria p. 201 (223) text-fig. 22, Chang Acta palaeont. sinica 7 3 1959.

Peishania lubrica sp. n. [nom. nud. 1957] pl. iv figs. 7-11 [sic], P. l. var. peitalingensis var. n. pl. iv fig. 6 p. 210 (234) Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Peltocare modestum sp. n. p. 158 pl. i figs. 9–10 Ordovician (Tremadoc) Norway, Henningsmoen Norsk geol. Tidsskr. 39 2–3 1959.

Perischodory Raymond 1937 considered a subgenus of Leiostegium [quod vide] p. 114, Berg & Ross.

Peronopsis fallax (Linnrs.) p. 5, Lazarenko.

Peronopsis interstrictus (White) p. 84 pl. xviii figs. 3-4c (sections), Bright.

Peronopsis cf. scutalis p. 12, P. cf. fallax p. 16, Kindle & Whittington.

Perunaspis Přibyl 1949 discussed and referred to Ceratarginae p. 138, P. longispina P. 1949 p. 139 pl. vii figs. 4-7 text-figs. 30-31, P. minuta (Barrande 1846) p. 141 pl. vii figs. 1-3 text-figs. 32-34, Vaněk (2).

Phacopacea see Phacopina subord. n.

Phacopacea—Ogygia sp. v. Freyberg 1923 may belong p. 41, Struve (1),

Phacopid new species [distinguished from Phacops rana (Green)] p. 1727, Jones.

Phacopida see Phacopina subord. n.

Phacopidae diagnosis of R. & E. Richter 1959 emended to accommodate *Volkops* gen. n. [quod vide] p. 32, Struve (1).

Phacopidae—phylogeny of Thuringian U. Devonian forms table 1 (p. 265), Pieiffer.

Phacopidella (Dienstina) liopyga Reinh. Richter neotype chosen and figured p. 271 pl. iv fig. 6, Pfeiffer.

Phacopidella (Denckmanniles) micromma (Roemer 1854) p. 71 2 text-figs., Strnad (6).

Phacopidina apiculata (M'Coy) so attributed and referred p. 146, Dean (2).

Phacopidina apiculata (M'Coy) p. 214, Dean (1).

Phacopina subord. n. [Phacopida] p. 461 for Phacopacea and Dalmanitacea, Struve Treatise on Invertebrate Paleontology Part O 1959.

Phacopinae see Volkops gen. n.

Phacops (Pterygometopus) dagon Reed may belong to new genus p. 147, Dean (2).

Phacops (P.) granulatus Mstr p. 269 pl. iii fig. l, P. (P.) cryphoides R. & E. Richter p. 269 pl. iii figs. 2-4, P. (Cryphops?) wocklumeriae R. & E. R. p. 270 pl. iii figs. 5a-b, 6, P. (Trimerocephalus) lacunosus sp. n. p. 270 pl. iv figs. 1-4, P. (T.) mastophthalmus Reinh. R. pl. iv fig. 5 Devonian (Upper) Germany (Thuringia), Pfeiffer Geologie 8 3 1959.

Phacops (Phacops) granulatus (Münster) p. 105, P. (Dianops) anophthalmus (Frech) p. 106 pl. ii fig. 3, Lewowicki,

Phacops (P.) granulatus (Münster), P. (Cryphops?) wocklumeriae R. & E. Richter p. 58, **Dearman &** 

Phacops latifrons—forged individual with 12 thoracic segments text-fig. 3, Schmidt.

Phacops (Dalmanites) longicaudatus var. orientalis Reed p. 1548, Bordet et al.

Phacops quadrilineatus Angelin pl. v fig. 19, Balashova (1).

Phacops rana (Green) see Phacopid new species.

Phacops (P.) sp. near sternbergi Corda p. 102 [for Ferronnière 1920–22 pl. 1 fig. 4], Pillet (2).

Phacops sp. p. 339ff., Paulus.

Phalacroma cf. glandiforme p. 17, Kindle & Whittington.

Pharostoma pediloba (Roem.) p. 6ff., Männil.

Pharostoma pulchrum pulchrum (Barr.) p. 48, Vaněk (3).

Phillipsia sp. aff. P. eichwaldi (Fisch.) p. 143, Ivanova.

Phillipsia scabra Woodward see Pygidia.

Phillipsia sp. nov. Koriyama p. 207, Yanagisawa.

Phillipsinella parabola (Barrande) p. 218, Dean (1).

Phorocephala similis n. gen. n. sp. [nom. nud.] p. 73 Ordovician China, Lu. Phylacops bituberculatus sp. n. (with transitory pygidia) including Aeglina rediviva of Baily 1862 p. 377 pl. lxii figs. 1-4 Ordovician (Upper) Eire, Weir Palaeontology 1 4 1959.

Phylacops recurvus n. sp. [nom. nud.] p. 77 Ordovician China, Lu.

Piriproetus see Parapalpebralia.

Placoparia cambrensis Hicks p. 227, Shackleton.

Planaspis Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Platycalymene tenera (Barr.) for Calymene t. B. 1852 p. 215, Vaněk (1).

Platylichas Gürich 1901 discussed and referred to Lichainae p. 114, P. kloučeki (Růžička 1926) lectotype chosen and figured including Lichas sp. R. 1926 (pl. 3 figs. 4-5) p. 115 pl. i fig. 3, Vaněk (2).

Platylichas laxatus (McCoy) p. 240, Shackleton.

Platylichas laxatus (M'Coy) s.l. p. 207ff., P. ef. l. p. 214, **Dean** (1).

Platypeltoides incipiens (Brøgger) listed p. 174, Henningsmoen.

Plethometopus armatus (Billings) p. 383 pl. lii fig. 14, pl. liii figs. 1–8, P. obtusus Rasetti including Unassigned pygidium no. 7 of R. 1945 p. 383 pl. liii figs. 11–14, P. convergens (Raymond) p. 384 pl. liii figs. 15–19, P. sp. undet. no. 1 p. 384 pl. liii figs. 9–10, P. sp. undet. no. 2 p. 384 pl. lii figs. 1–4; Unassigned pygidia nos. 9 and 10 of Rasetti 1945 belong to P. (possibly to P. a.) p. 383, Rasetti.

Plethopeltella see Iranaspidinae subfam. n.

Plicatolina sp. from Norway p. 174, Henningsmoen.

Pliomera fischeri (Eichwald) 1825 p. 311 pl. lxxxiv figs. 1–3b, P. brevicapitata (Lamansky) 1905 pl. lxxi figs. 4a–5, Lesnikova & Weber.

Pliomeridae see Diaphanometopinae subfam. n.

Pliomeroides see Protopliomeroides.

Pliomeroides primigenus (Angelin) listed p. 174, Henningsmoen.

Poliellaspis Lermontova discussed p. 113, P. sayanicus (Poletaeva) lectotype selected [illegitimate] p. 114 pl. v figs. 14–15, Pokrovskaya.

Poliellaspis range in Altai-Sayan pl. i, Zhuravleva & Repina.

Poliellaspis sp. p. 1033, Zhuravleva et al. (2).

Poliellina Poletaeva discussed p. 100, P. lermontovae P. lectotype selected [illegitimate] p. 102 pl. v figs. 1, 4, 7–8, 10, 13, P. elongata sp. n. p. 104 pl. v figs. 2–3, 5–6, P. cylindrica sp. n. p. 107 pl. v figs. 9, 11–12, P. sicropyge sp. n. p. 109 pl. vi figs. 10–11 Cambrian (Lower) U.S.S.R. (Tuva etc.); P. sp. [det. Poletaeva] p. 16, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959

Poliellina range in Altai-Sayan pl. i, Zhuravleva & Repina.

Pr Pr

[19

Po

type (223)

pl. ii

(224)

Acta

Prince Pr

Bla [non 3-4 [Re visi sini

E. 1

pale

Re

Tu

P

19

p

fi s

Poshania gen. n. [Namanoiidae] p. 200 (221) for type P. poshanensis sp. n. [nom. nud. 1957] p. 201 (223) pl. ii figs. 4-10 text-fig. 21, P. transversa sp. n. pl. ii figs. 11-13 text-fig. 20, P. sp. pl. ii fig. 14 p. 203 (224) Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Primaspis ascitus p. 388, Whittington.

Primaspis keyserlingi (Barr.) p. 48, Vaněk (3).

Primaspis semievoluta (Reed) p. 214, Dean (1).

Proasaphiscus yabei Resser & Endo stated to have 11 thoracic segs p. 226 text-fig. 23, P. (Honanaspis) subgen. n. p. 204 (226) for type P. (H.) honanensis sp. n. p. 205 (227) pl. iii figs. 1–4 text-fig. 24, P. (H.) lui sp. n. [nom. nud. 1957] p. 206 (228) pl. iii figs. 5–6 text-fig. 25 Cambrian (Middle) China (Honan & Shantung), P. (H.) machidai (Endo) so referred p. 229; Manchuriella R. & E. probably a subgenus of P. p. 227, Conokephalina termieri and tienfongensis Mansuy 1916 represent a new subgenus of P. p. 227n, P. centronatus R. & E. and tentatively P. offula R. & E. referred to Szeaspis gen. n. p. 230, Chang Acta palaeont. sinica 7 3 1959.

Prodamesella gen. n. [compared with Damesella and Blackwelderia] p. 196 (218) for type P. convexa sp. n. [nom. nud. 1957] p. 196 (219) pl. i fig. 13 text-figs. 3-4 Cambrian (Middle) China (Shantung), D. quadrata [Resser &] Endo & Olenoides manchuriensis E. provisionally referred to P. p. 219, Chang Acta palaeont. sinica 7 3 1959.

Proceio Lerm, discussed, P. quadratica sp. n. p. 146 pl. vii figs. 1-6, 8-9 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Proerbiarange in Altai-Sayan pl. i, Zhuravleva & Repina.

Proerbia torgaschinica Repina [nom. nud.] p. 1032, Zhuravleva et al. (2).

Proetacea see Illaenina subord, n.

Proetid p. 1039, Holland et al.

Proetidarum gen. et sp. indet. p. 349, Paulus.

Proetidella aff, fearnsidesi Bancroft p. 206, Dean (1).

Proetids p. 384ff., Whittington.

Proetus cultrijugati ? R. & E. Richter, P. sp. p. 339, P. sp. Gruppe c. p. 349, **Paulus**.

Proetus (P.) cuvieri Steininger [for Ferronnière 1920-22 pl. 1 fig. 10] p. 102, P. (P.) orbitatus Barrande p. 103, Pillet (2).

Proetus ? primulus Barrande referred to Ogygites p. 46, Vaněk (3).

Proetus ramisulcatus Nieszkowski 1857 p. 29 pl. i fig. 3, Balashova (1).

Proetus sp. cf. P. regalus Poulsen p. 50, Thorsteinsson.

Proetus romanovskii Weber 1932 p. 299 pl. lxxix figs. 5a-b, Lesnikova & Weber.

" Proetus" strasburgensis p. 384, Whittington.

Proetus sp. p. 514, Walmsley.

Proliostracus strenuelliformis Poulsen 1932—concluded from plots of dimensions that P. rosenkrantzi, P. liostracoides and P. noe-nygaardi P. are synonyms p. 474ff. text-figs. 1-15, Shaw.

Promegalaspides see Promegalaspidinae subfam, n.

Promegalaspides intactus (Moberg & Segerberg) p. 274, Tjernvik.

Promegalaspides (Borogothus) intactus (Mob. & Seg.) listed p. 174, Henningsmoen,

Promegalaspides kasachstanensis Bal. [not seen] p. 144, Balashova & Balashov.

Promegalaspidinae subfam. n. [Asaphidae] p. 352 for *Promegalaspides* and *Borogothus*, **Jaanusson** Treatise on Invertebrate Paleontology Part O 1959.

Prosaukia stosei (Walcott) supplementary description p. 389 pl. liv figs. 16–22, P. corrugata sp. n. with immature stages p. 390 pl. liv figs. 4–14 Cambrian (Upper) U.S.A. (Pa & Va), Rasetti J. Paleont. 33 3 1959.

Protolenidae discussed p. 395, relations to Dolerolenidae and Lermontoviidae p. 396, **Sdzuy**.

Protoleninae Hupé discussed with synonym Lermontoviinae Suvorova p. 39 [see also Elegestina and Kadyella genn. n.], Pokrovskaya.

Protolenus Matthew 1892 discussed with synonyms Matthewlenus Hupé & Lermontovia Suvorova p. 42, P. planus 5p. n. p. 43 pl. ii figs. 11, 14, 18 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Protolenus n. sp. [undescr.] pl. i fig. 7, Sdzuy.

Protopliomeroides [altered in ink to Pliomeroides in offprint from author] primigenius (Angelin) p. 274, Tjernvik.

Protopliomerops speciosus Dalm., P. sp. p. 144, Balashova & Balashov.

Protypus sp. p. 1032, Zhuravleva et al. (2).

Pruvostina Hupé 1952 [sic] referred to Bigotininae p. 401, Sdzuy.

Pruvostinoides Hupé 1952 [sic] referred tentatively to Bigotininae p. 401, Sdzuy.

Psalikilus—Gonioteloides may be related p. 117, P. paraspinosum Hintze pygidium illustrated pl. xxi fig. 5, Berg & Ross.

Pseudagnostus convergens Palmer p. 412 pl. lvii figs. 1-6, P. prolongus (Hall & Whitfield) with muscle scars p. 412 pl. lvii figs. 7-16 text-fig. 1, Lochman & Hu.

Pseudagnostus of. cyclopyge p. 15, Kindle & Whittington.

Pseudagnostus gyps (Clark) p. 381 pl. li figs. 13-14, Rasetti

Pscudasaphus globifrons (Eichwald) 1857 p. 288 pl. lxxiv figs. 1-2, P. tecticaudatus (Steinhard) 1874 p. 289 pl. lxxiii fig. 1, Lesnikova & Weber.

Pseudasaphus huangi n. sp. [nom. nud.] p. 76 Ordovician China, Lu.

[19

p. 40

pl. iv

figs.

pl. 2 figs.

pl. :

stag

R. e

p. 4

wh

195

p. 2

gin

figs

P. 184

gro

fig W

R

St

1

Pseudoeteraspis sp. p. 1372, Z. A. Zhuravleva (2).

Pseudogygites latimarginatus (Hall) p. 46ff., horizon discussed p. 89, P. sp. p. 89n, Thorsteinsson.

Pseudoolenoides ? recorded p. 111, Berg & Ross.

Pseudosphaerexchus [sic] orvikui Männ. pl. xxix fig. 4, Kalio (2).

Pseudosphaerexochus cf. granulatus Ang. p. 8, Männil.

Pseudosphaerexochus octolobatus p. 204, P. sp. nov. [undescr.] p. 207ff., **Dean** (1).

Pseudosphaerexochus pahnschi Schmidt 1881 p. 309 pl. lxxxiii figs. 11-12, Lesnikova & Weber.

Pseudosphaerexochus sp. p. 379 pl. lxii fig. 16, Weir.

Psilocephalina lata n. sp. [nom. nud.] p. 36 Ordovician (Lower) China, Lu.

Ptarmigania sp. undet. p. 195 pl. xix fig. 33, Resser.

Pterocephalia recorded p. 107, J. BRIDGE in Berg & Ross.

Pterocephaliidae referred to Olenina and probably closely related to Olenidae p. 157, Henningsmoen.

Pterygometopus see Phacops.

Pterygometopus huayinshanensis n. sp. [nom. nud.] p. 43 Ordovician China, Lu.

Pterygometopus jukesi (Salter) p. 240, Shackleton.

Pterygometopus nieszkowskii (Schmidt) 1881 p. 315 pl. lxxxiv figs. 9a-10, P. sclerops (Dalm.) 1826 p. 315 pl. lxxxiv figs. 7-8, Lesnikova & Weber.

Pterygometopus sp. Bancroft 1948 probably belongs to Duftonia lacunosa gen. et sp. n. p. 146, P. jukesi (Salter) referred to Estoniops, Dean (2).

Ptychagnostus showing genal caeca text-fig. 1, Önik (1),

Ptychagnostus atavus p. 17, P. ef. a. p. 12, P. ef. aculeatus p. 17, Kindle & Whittington.

Ptychaspididae see Olenina.

Ptychaspis bullasa sp. n. with immature stages p. 422 pl. lviii figs. 21-42, P. granulosa (Owen) p. 424 pl. lx figs. 37-44 Cambrian (Upper) U.S.A. (Idaho), Lochman & Hu J. Paleont. 33 3 1959.

Ptychaspis granulosa (Owen) p. 112 pl. xxii fig. 12; pygidium possibly of P. p. 107, Berg & Ross.

Ptychoparia clusia Walc. p. 30; P. ? sp. Poletaeva 1936 referred to Lermontoviella sayanica gen. et sp. n. p. 155, Pokrovskaya.

Ptychoparia truncata Sampelayo p. 113 (pl. 39) fig., C. fichti Walcott p. 114 (pl. 40) fig., C. azpeitiae S. p. 115 (pl. 41) fig., Badillo.

Ptychoparia sp. p. 5, Lazarenko.

Ptychopariida see Harpina & Illaenina subordd, n.

Ptychopariids—traces of intestinal appendages on thorax and pygidium p. 1751, Öpik (1).

Ptychopyge Angelin 1854 placed on Official List of Generic Names and type designated Asaphus angustifrons Dalman 1827 [quod vide] p. 59, Opinion 538.

Ptychopyge angustifrons (Dalman) 1827 p. 288 pl. lxxii fig. 5, Lesnikova & Weber.

Ptychopyge? inostranzewi Lamansky 1905 referred to Asaphellus p. 145, Balashova & Balashov.

Pumilina Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Raphiophoridae—discussion of morphology and ontogeny of Ampyx, Lonchodomas, Ampyxina, Raymondella p. 460ff., Whittington.

Raphiophoridae see Ampyxinellinae subfam. n.

Raphiophorus volborthi Schmidt 1894 p. 276 pl. lxv figs. 5, 7, R. bulbifer Weber 1932 p. 276 pl. lxv fig. 8, Lesnikova & Weber.

Rawlinsella see Bolaspididae fam. n.

Raymondaspis gregarius p. 384ff., Whittington.

Raymondella Reed 1935 discussed p. 487, R. elegans (Cooper 1953) for Ampyxina e. C. restr. [non pl. 4 figs. 10-11, 13, 17] p. 488 pl. xxxvi figs. 1-37, R. sp. for A. sp. C. 1953 p. 490 pl. xxviii figs. 15-16, 19, Whittington.

Raymondella nordica sp. nov. pl. vi figs. 15-16, Balashova (1).

Redlichiacea see Despujolsiidae & Gigantopygidae fam, n.

Redlichiacea see Redlichiina subord. n.

Redlichiida see Redlichiina & Bathynotina subordd. n.

Redlichiidae see Asthenaspis gen. n.

Redlichiina subord. n. [Redlichiida] p. 198 for Redlichiacea, Ellipsocephalacea and Paradoxidacea, Harrington Treatise on Invertebrate Paleontology Part O 1959.

Redlichina Lermontova 1940 discussed p. 65, R. lermontovae sp. n. p. 68 pl. iv figs. 8-11, R. rarissima sp. n. p. 70 pl. iv figs. 2, 4, 6, R. tuberculata sp. n. p. 72 pl. iii figs. 11-14, 16, R. angusta sp. n. p. 76 pl. iii fig. 15, R. pustulosa sp. n. p. 79 pl. iv figs. 1, 3, 5, R. dubia sp. n. p. 81 pl. iv fig. 7 Cambrian (Lower) U.S.S.R. (Tuva), R. zabaikalica [nom. nud.] p. 67, R. sp. p. 29, Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Redlichina Lermontova 1940 referred tentatively to Doleroleninae p. 400, Sdzuy.

Redlichoidea see Menneraspidae fam. n.

Reedolithus nováki (Chlupáč) for Tretaspis n. C. 1952 with which Trinucleus alfredi Želízko is probably synonymous p. 215, Vaněk (1).

Reedolithus nováki (Chlupáč) p. 48, Vaněk (3).

Remopleuridacea see Loganellidae fam. n.

Remopleurides Portlock 1843 diagnosis with discussion of morphology (including comparison of hypostome with that of Ceraurus) and ontogeny (including protaspides) p. 391ff., R. caelatus 5D. R.

p. 401 pl. i figs. 1–16, pl. ii figs. 1–25, pl. iii figs. 1–18, pl. iv figs. 1–25, text-figs. 3c, 4a-c, 5a-B, R. plaesiourus sp. n. p. 412 pl. iv figs. 26–30, pl. v figs. 1–26, pl. vi figs. 1–15, R. caphyroides sp. n. p. 414 pl. vii figs. 1–10, pl. viii figs. 1–10, pl. ix figs. 1–22, pl. x figs. 1–20, pl. xii figs. 1–4, text-figs. 6a-B, R. c. ? p. 419 pl. xi figs. 5–11, R. asperulus sp. n. p. 420 pl. xii figs. 1–20, pl. xiii figs. 1–31, R. eximius sp. n. p. 424 pl. xiv figs. 1–31, R. eximius sp. n. p. 424 pl. xiv figs. 1–4, pl. xv figs. 1–18, R. e. ? [developmental stages tentatively referred] p. 425 pl. xvi figs. 1–17, R. simulus sp. n. [may be sexual dimorphs of R. e.] p. 426 pl. xvi figs. 1–23, pl. xix figs. 11–12 Ordovician (Middle) U.S.A. (Virginia), R. rugicostatus Raymond type refigured p. 427 pl. xviii figs. 23–24, 26–27, Whittington Bull. Mus. comp. Zool. Harv. 121 8

Remopleurides aff. burmeisteri Bancroft, R. sp. p. 207, R. sp. ind. p. 208, Dean (1).

Remopleurides of. dalecarlicus Warb. p. 7, R. emarginata Törnq. p. 11, Männil.

Remopleurides pisiformis Weber 1948 p. 277 pl. lxvi figs. 1a-c, 2, Lesnikova & Weber.

Remopleurides cf. salteri var. girvanensis Reed 1903 p. 26 pl. i figs. 4a-5, R. aff. longicostatus Portlock 1843 p. 27 pl. i fig. 2, Balashova (1).

Remopleurides sp. p. 373 pl. lxii fig. 20, Weir.

Remopleuridid gen. et sp. ind. [may be early growth-stages of *Robergiella eagittalis*] p. 434 pl. xix figs. `1-10 Ordovician (Middle) U.S.A. (Virginia), Whittington.

Remopleurididae see Olenina.

Remopleuridinae see Robergiella gen. n.

Resimopsis range in Altai-Sayan pl. i, Zhuravleva & Renina.

Richardsonella near R. invita (Salter) p. 56, Stubblefield (1).

Richardsonella megalops (Billings) p. 386 pl. lv fig. 14, R. subcristata sp. n. p. 386 pl. lv figs. 15–18, R. cf. unisulcata Rasetti p. 387 pl. lv fig. 19 Cambrian (Upper) U.S.A. (Maryland), Rasetti J. Paleont. 33 3 1959.

Robergia barrandei (Etheridge et Nicholson) 1879 p. 25 pl. i figs. 18-19, Balashova (1).

Robergia major Raymond 1920 with synonym R. athenia Butts 1926 p. 428 pl. xviii figs. 1-22, 25, Whittington.

Robergiella gen. n. [Remopleuridinae] p. 431 for type R. sagittalis sp. n. p. 432 pl. vi figs. 16-33 [? figs. 20-22 p. 433—see also Remopleuridid gen. et sp. ind.] Ordovician (Middle) U.S.A. (Virginia), Whittington Bull. Mus. comp. Zool. Harv. 121 8 1959.

Rondocephalus gen. n. [Dolichometopinae] p. 122 for type R. mirandus sp. n. p. 123 pl. vii figs. 11-17 Cambrian (Lower) U.S.S.R. (Tuva etc.), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Rondocephalus range in Altai-Sayan pl. i, Zhurav-leva & Repina.

Sajanaspis Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Saltaspis recorded from Britain p. 155, Stubblefield (3).

Salteria americana Cooper 1953 p. 491 pl. xxviii figs. 9, 12, 14, 17, 18, Whittington.

Sanashtycgolia [Ianashtycgolia on pl.] Poletaeva [nom. nud.] range in Altai-Sayan p. 182 pl. i, Zhuravleva & Repina.

Saratogia Walcott 1916 discussed with diagnosis and relations to Idahoia p. 421, S. fria sp. n. p. 422 pl. lix figs. 1-11 Cambrian (Upper) U.S.A. (Idaho), Lochman & Hu J. Palcont. 33 3 1959.

Saukianda andalusiae R. & E. Richter 1940 pygidium with posteriorly incomplete doublure (considered not a sexually dimorphic character) p. 403 text-fig. 1, Sdzuy.

Saukiid (undetermined) p. 379, Rasetti.

Saukiidae see Olenina,

Sayanella gen. n. [Dinesidae] p. 150 for type Klotziella poletaevae [poletayevi] Lermontova 1940 lectotype selected [illegitimate] p. 152 pl. ix figs. 14–15 Cambrian (Lower) U.S.S.R. (Sayan & Tuva), S. bona [nom. nud.] p. 151, Pokrovskaya Trud. geol. Inst. S.S.R. 27 1959.

Schmidtella see Cyphaspis.

Schmidtiellus mickwitzi (Schm.) [reconstruction] text-fig. 3 (1), Kaljo (1).

Scutelluidae see Goldillaeninae subfam. n.

Scutelluidae see Thysanopeltidae.

Scutellum see Bronteus.

Scutellum costatum Pusch p. 269 pl. i fig. 5, Pfeiffer,

Scutellum taimyricum sp. nov. pl. i figs. 10, 10a, Balashova (1).

Scutellum taimyricum sp. n. p. 28 [pl. i figs. 10, 10a] Silurian (Lower) U.S.S.R. (Arctic), Balashova Sborn, stat. paleont, biostrat. 15 [pls. in 14] 1959.

 $Scutellum\left\{S.\right\}$ n. sp. [undescr.] p. 102 [for Ferronnière 1920–22 pl. 1 figs. 1–2], S. (S.)3 spp., S. (Paralejulus) sp. p. 103, Pillet (2).

Scutellum sp. p. 353, Paulus.

Selenopeltis buchi buchi (Barr.) p. 48, Vaněk (3).

Serrodiscus R. & E. Richter discussed p. 170, S. cf. speciosus (Ford) p. 172 pl. xi fig. 1, S. sibiricus sp. n. p. 173 pl. xi figs. 2-4, 9, 21, S. f granulatus sp. n. p. 176 pl. xi fig. 18 Cambrian (Lower) U.S.S.R. (Tuva) [S. communis p. 35 and S. asiaticus p. 171 are apparently misprints for S. sibiricus], Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Shanganella gen. n. [Incertae sedis] p. 161 for type S. lebedevae sp. n. p. 162 pl. xi figs. 22-23 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Shivelicus gen. n. [Pagetiidae] p. 180 for type S. parvus gp. n. p. 181 pl. x figs. 3, 7, 9-10, 12-13, 15-16 Cambrian (Lower) U.S.S.R. (Tuva, Sayan etc.), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Shivelicus range in Altai-Sayan pl. i, Zhuravleva & Repina.

Shumardia extensa sp. n. p. 374 pl. lxiii figs. 5-6 Ordovician (Upper) Eire, Weir Palaeontology 1 4 1959.

Shumardia lata n. sp. p. 36, S. gonioloba n. sp. p. 38 [nom. nuda] Ordovician China, Lu.

Shumardia oelandica Moberg p. 144, Balashova & Balashov.

Shumardia pusilla (Sars) listed p. 174, Hennings-moen.

Shumardia cf. pusilla (Sars) p. 274, Tjernvik.

Skemmatopyge tietzei R. & E. Richter p. 105, Lewowicki.

Skemmatopyge tietzei R. & E. Richter p. 268 pl. i fig. 4, Pfeiffer.

Solenopleura sp. p. 359, Miroshnikov et al.

Solenopleurella bella Lerm. p. 798, Z. A. Zhuravleva (1).

Solenopleurella porcata sp. n. p. 214 pl. xxvii figs. 33-34, S. erosa sp. n. p. 214 pl. xxiii figs. 5-6, pl. xxvii figs. 26-28, S. diligens sp. n. p. 215 pl. xxvii figs. 29-31, S. sp. undet. p. 215 pl. xxvii fig. 32 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg, Instn No. 563 1945.

Solenopleuropsis marginata angularis Sdzuy p. 116 (pl. 42) fig., Badillo.

Solontzella Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Spencia tontoensis sp. n. p. 213 pl. xxiv figs. 23-30, pl. xxv figs. 1-2 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Sphaerexochus angustifrons Ang. p. 11, Männil.

Sphaerexochus hapsidotus p. 384, S. pulcher p. 385, Whittington.

Sphaerexochus hisingeri Warburg 1925 p. 308 pl. lxxxiv figs. 5a-b, S. mirus Beyrich 1845 p. 308 pl. lxxxiv fig. 6, Lesnikova & Weber.

Sphaerexochus taimyricus sp. nov. pl. v figs. 8a-6, Balashova (1).

Sphaerexochus sp. p. 1548, Bordet et al.

Sphaerocoryphe hübneri Schmidt 1881 p. 311 pl. lxxxii fig. 12, Lesnikova & Weber.

Sphaerocoryphe cf. thomsoni p. 204, Dean (1).

Staurocephalus aff. murchisoni Barrande including S. globiceps of Baily 1862 p. 379 pl. lxiii figs. 2-3, Weir.

Staurocephalus sp. Weber 1948 referred to Ovalocephalus kelleri gen. et sp. n. p. 1316, Koroleva.

Stenochilina spinifera Ulrich p. 383 pl. lii figs. 15–16, Rasetti.

Stenopareia avus Holm pl. iii figs. 1-4, S. borealica sp. nov. pl. iii figs. 15-19, Balashova (1),

Stenopareia avus Holm p. 42 [pl. iii figs. 1-4], S. borealica sp. n. Ordovician (Upper) U.S.S.R. (Arctic) p. 43 [pl. iii figs. 15-19], Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Stenopareia panderi (Barr.) p. 47, Vaněk (3).

Stenopilus Clark 1924 so accredited with type S. intermedius C. p. 385, S. pronus Raymond [not type of S.] p. 385 pl. liii figs. 20-24, S. sp. undet. p. 377, S. sp. p. 379, Rasetti.

Strenuaeva primaeva (Brögger) p. 521 pl. i figs. 1-9, S. orlovinensis (Czarnocki) [sp. n.] neotype [sic] for Strenuella o. C. 1927 nom. nud. p. 522 pl. i figs. 10-15, S. kiaeri (C.) [sp. n.] for Strenuella k. C. 1927 nom. nud. p. 523 pl. i figs. 16a-c Cambrian (Lower) Poland (H. Cross Mts.), Samsonowicz Bull. Acad. polon. Sci. [sci. chim., etc.] 7 7 1959.

Strenuaeva primaeva pl. i fig. 4 pars, Samsonowicz (3).

Strenuella tentatively recorded p. 771, Orlowski (1).

Strenuella orlowinensis and S. kiaeri Czarnocki see Strenuaeva.

Strenuella polonica Czarnocki nom. nud. 1927 [5p. n.] neotype [sic] p. 525 pl. i figs. 1–12, S. cf. lakei Hupé p. 526 pl. ii figs. 1–3 Cambrian (Lower) Poland (H. Cross Mts.), Samsonowicz Bull. Acad. polon. Sci. [sci. chim., etc.] 7 7 1959.

Stygina latifrone Portlock pl. ii figs. 11-12, Balashova (1).

Stygina latifrons (Portlock 1843) p. 27, Balashova (2).

Symphysops aff. subarmatus (Reed) p. 378 pl. lxiii figs. 10-11, Weir.

Symphysurina sp. cf. S. globocapitella Hintze p. 112 pl. xxi fig. 25, pl. xxii fig. 3, Berg & Ross.

Symphysurus angustatus (Sars & Boeck) listed p. 174, Henningsmoen.

Symphysurus angustatus (Sars & Boeck) p. 274, Tjernvik.

Symphysurus carinatu, S. subquadratus p. 76, S. nanshanensis p. 88 n. spp. [nom. nuda] Ordovician China, Lu.

Symphysurus incipiens Brögg., S. convexus Bal. [not seen] p. 144, Balashova & Balashov.

Synhomalonotus expansus n. sp. p. 74, S. concavus var. hous n. var. p. 76 [nom. nuda] Ordovician China, Lu.

Synphoria Clarke 1900 considered valid and so dated, Eccorycephalus Reed 1925 and Necephphoria Pillet 1954 objective synonyms p. 472n, STRUVE in Harrington et al.

Syspacephalus asperoensis Rusc. discussed p. 9, Rusconi.

Szeaspie gen. n. [no family assigned, compared with i. a. Asaphiscus and Anomocare] p. 207 (229) for typo S. reticulatus sp. n. for Asaphiscus cf. iddingsi of Chang 1957 pl. iii figs. 11-16, S. r. var. brevicus var. n. pl. iii figs. 7-10 p. 208 (230) Cambrian (Middle) China (Shantung), S. pustulosa (Endo & Resser) for

Manch S. cent S. ? of (Walco referre fig. 4 p

Γ195

7ae 34–43 Hu J.

Szec

Tae cranid Minke

Tai rica \$1 U.S.S Sborn

> spp. (Midd 17–18 T. gr p. 22 R. & Acta

with

Trud type (Upp Sci. 1

Camb

Te figs.

Te

Ordo

T) 11-1 T, g pl. h

Bron 1955

pl. 1
Tr

T

for Jaar Pari Manchuriella p. R. & E. 1937 restr. (pl. 36 fig. 13), S. centronatus (E. & R.) for Proasophisous c. R. & E., S. ? offula (E. & R.) for P. o. R. & E., S. iddingsi (Walcott) for Asaphiscus i. W., S. sp. for free cheek referred to Fuchouia quadrata by R. & E. 1937 (pl. 39 fig. 4 pars) p. 230, Chang Acta palaeont. sinica 7 3 1959.

Szechuanella szechuanensis n. gen. n. sp. [nom. nud.] p. 33 Ordovician (Lower) China, Lu.

Taenicephalina globula sp. n. p. 424 pl. lvii figs. 34–43 Cambrian (Upper) U.S.A. (Idaho), Lochman & Hu J. Paleont. 33 3 1959.

Taenicephalus megalops Kobayashi—some referred cranidia of K. and of Wilson 1954 may belong to Minkella gen. n. p. 416, Lochman & Hu.

Taimyraspis gen, n. [Isocolidae] for type T. taimyrica sp. n. p. 19 pl. vi figs. 17-24 Ordovician (Middle) U.S.S.R. (Arctic) [see also Holdenia typa], Balashova Sborn. stat. paleont. biostrat. 14 1959.

Taitzuia Resser & Endo referred to Namanoiidae with probable synonym Inouyina p. 223, review of spp. of T. p. 225, T. quadrata sp. n. Cambrian (Middle) China (Shantung) p. 203 (224) pl. iv figs. 17–18; T. adrastia (Walcott) referred to Levisia, T. granulata Endo provisionally to a new genus p. 226; pygidium attributed to Aojia longispina R. & E. 1937 (pl. 47 fig. 2) belongs to T. p. 223, Chang Acta palaeont. sinica 7 3 1959.

Tannudiscus gen. n. [Opsidiscidae] p. 177 for type T. tannudiscus sp. n. p. 178 pl. xi figs. 16, 19-20 Cambrian (Lower) U.S.S.R. (Tuva), Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Tatulaspis gen. n. [no family assigned] p. 117 for type T. princeps sp. n. p. 118 pl. i figs. 6, 6a Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Telephina aff. reedi (Ulrich) p. 373 pl. lxiii figs. la-c, Weir.

Tetralichas holmi (Schmidt) 1885 p. 303 pl. lxxx figs. 3a-4, Lesnikova & Weber.

Tetralichas taimyricus sp. nov. pl. i figs. 5-6, Balashova (1).

Tetralichas taimyricus sp. n. p. 50 [pl. i figs. 5-6] Ordovician (Upper) U.S.S.R. (Arctie), Balashova Sborn. stat. paleont. biostrat. 15 [pls. in 14] 1959.

Theodenisia communis (Rasetti) p. 383 pl. li figs. 11-12, T. sp. undet. p. 380, Rasetti.

Thomondia Harper 1942 referred to Isocolidae,  $T.\ globosa\ H.$  thorax and pygidium described p. 375 pl. lxii figs. 11-15, **Weir**.

Thysanopeltidae Hawle & Corda 1847 preferred to Bronteidae H. & C. or Scutelluidae R. & E. Richter 1955 p. 367, Moore in Harrington et al.

Thysanopeltidae see Weberopeltis gen, n,

Thysanopeltis sp. p. 102 [for Ferronnière 1920-22 pl. 1 fig. 3], T. sp. near speciosa p. 103, Pillet (2).

Thysanopyge see Thysanopyginae subfam. n.

Thysanopyginae subfam. n. [Asaphidae] p. 354 for Thysanopyge, Australopyge and Zuninaspis, Jaanusson Treatise on Invertebrate Paleontology Part O 1959.

Tomagnostus cf. fissus p. 17, Kindle & Whittington.

Tonkinella shantungensis sp. n. [nom. nud. 1957] p. 208 (231) pl. iv fig. 1 Cambrian (Middle) China (Shantung), Chang Acta palaeont. sinica 7 3 1959.

Trachycheilus gen. n. [no family assigned, compared with Kochina] p. 212 for type T. typicale sp. n. p. 213 ppl. xxv fig. 9 pars Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Tretaspis ef. bucklandi (Barrande) 1846 p. 275 pl. lxxix figs. 6a-c, Lesnikova & Weber.

Tretaspis cf. ceriodes (Angelin) p. 207ff., T. aff. kiaeri Størmer p. 208, Dean (1).

Tretaspis nováki Chlupáč referred to Reedolithus [quod vide] p. 215, Vaněk (1).

Tretaspis sagenosus sp. n, including T. reticulatus of Whittington 1941 partim (pl. 6 figs. 30, 34, 35) & Ampyxina elegans Cooper 1953 partim (pl. 4 figs. 13, 17) morphology and ontogeny described p. 448 pl. xxiv figs. 1-19, pl. xxv figs. 1-6, pl. xxvi figs. 1-17, pl. xxvii figs. 1-25, pl. xxviii figs. 1-8 Ordovician (Middle) U.S.A. (Virginia), Whittington Bull. Mus. comp. Zool. Harv. 121 8 1959.

Triarthropsis nitida Ulrich p. 382 pl. lv figs. 6–13, T. limbata sp. n. Cambrian (Upper) U.S.A. (Maryland) p. 382 pl. lii figs. 1–8, Rasetti J. Paleont. 33 3 1959.

Triarthrus sp. aff. becki p. 89n, Thorsteinsson.

Trilobita Walch 1771—type of class designated Dalmanites Barrande 1852 p. 172, HARRINGTON, HENNINGSMOEN & MOORE in Harrington et al.

Trilobus punctatus Brünnich 1781 suppressed p. 43, Opinion 537.

Trimerocephalus Boo Phacops.

Trinodus elspethi p. 384ff., Whittington.

Trinodus glabratus var. kirgizica Weber 1948 p. 274 pl. lxv figs. 1a-c, Lesnikova & Weber.

Trinodus mobergi Tjernvik from Norway p. 174, Henningsmoen.

Trinodus ? sp. p. 371 pl. lxiii figs. 9a-b, Weir.

Trinucleidae—ontogeny discussed p. 439, Whittington.

Trinucleidae see Collis gen. n.

Trinucleoidea—morphology and ontogeny discussed with diagnosis p. 437, Whittington.

Trinucleus alfredi Želizko probably referable to Reedolithus and probably a synonym of R. nováki (Chlupáč) p. 215, Vaněk (1).

Trinucleus concentricus of Baily 1862 referred to Novaspis aff. albida (Reed) p. 371, Weir.

Triplagnostus of. hybridus p. 12, Kindle & Whittington.

Trochurus Beyrich 1845 discussed p. 144, T. speciosus B. 1845 neotype selected and figured [excluding record of Hawle & Corda 1847 and with synonyms after Barrande 1852 including tentatively Dicranopeltis? balkhaschicus and D. ? turensis Weber 1951] p. 145 pl. viii fig. 10, pl. ix figs. 1–9, pl. x figs. 1–7, pl. xi figs. 1–3, text-figs. 35–38, Vaněk (2).

dur

inde

to

Tropidopyge Harrington & Kay 1951 discussed and family reference uncertain p. 160, T. broeggeri (Moberg & Segerberg 1906) p. 159 pl. i figs. 5-7, T. stenorhachis H. & K. may not be congeneric p. 160, Henningsmoen.

Tropydocoryphe sp. [for Ferronnière 1920–22 pl. 1 fig. 9] p. 102, Pillet (2).

Tumulina Repina [nom. nud.] range in Altai-Sayan pl. i, Zhuravleva & Repina.

Tuvaaspis minutus [nom. nud.] p. 28 apparently a misprint for Tuvanella gracitis gen. et sp. n. [quod vide], Pokrovskaya.

Tuvanella gen. n. [Aldonaidae] for type T. gracilis sp. n. p. 95 pl. viii figs. 7, 11 Cambrian (Lower) U.S.S.R. (Tuva) [Tuvaaspis minutus p. 28 is apparently a misprint for Tuvanella gracilis], Pokrovskaya Trud. geol. Inst. S.S.S.R. 27 1959.

Tuvanella range in Altai-Sayan pl. i, Zhuravleva & Repina.

Tuvinia tannu-olae nov. gen. sp. nov. [nom. nud.] [Corynexochidae] p. 16, O. K. POLETAEVA in Pokrovskaya.

Typhloproetus hemisphaericus sp. n. p. 266 pl. i figs. 1-2 Devonian (Upper) Germany (Thuringia), Pfeiffer Geologie 8 3 1959.

Typhloproetus subcarintiacus R. & E. Richter p. 104, Lewowicki.

Ulrichaspis text-fig. 3b, Kindle & Whittington.

Uralichas Delgado 1892 discussed p. 101, U. avus (Barrande 1872) restr. [non pl. 5 fig. 23, pl. 6 figs. 25–26, pl. 10 fig. 17 to Lichas incola] lectotype chosen and figured excluding record of G. & H. Termier 1950 and with synonym Autoloxolichas reedi Phleger p. 102 pl. ii figs. 1–6 text-figs. 6–9, U. giganteus (Klouček 1916) nov. emend. lectotype chosen and figured p. 107 pl. i figs. 1–2, Vaněk (2).

Urbanaspis gen. n. [no family assigned] p. 118 for type U. notabilis sp. n. p. 119 pl. i figs. 9, 9a Cambrian (Upper) U.S.S.R. (Kazakhstan), Ivshin Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955.

Varvia longicauda Tjernvik from Norway p. 174, Henningsmoen.

Viñakainella see Oryctocephalus.

Vogdesia nordica sp. n. p. 37 pl. ii figs. 5-6 Ordovician (Middle) U.S.S.R. (Arotic), Balashova Sborn, stat. paleont. biostrat. 14 1959.

Volkops gen. n. [Phacopinae] p. 31 for type V.volki sp. n. with genal and pleural spines Ordovician Germany (Thuringia) p. 35 pl. i figs. 1a-i, k, pl. ii figs. 1l-q, 2a-d (reconstructions), Struve Senckenbergiana leth. 40 1-2 1959.

Waribole see Cyrtosymbole.

Weberides mucronatus (M°Coy 1844) from Czechoslovakia considered to show sexual dimorphism p. 255 pl. i figs. 1-4, pl. ii figs. 1-4, pl. iii figs. 1-4, pl. iii figs. 1-4, pl. iv figs. 1-4 text-figs. 1-2, 4, Rehof & Rehořová.

Weberopeltis Maksimova 1957 [gen. n.] [Thysanopeltidae] p. 372 for type Bronteus aculeatus Weber 1945 text-fig. 281, R. & E. Richter Treatise on Invertebrate Paleontology Part O 1959.

Wilbernia pero (Walcott) p. 422 pl. lx figs. 15-18, Lochman & Hu.

Wutingaspis Kobayashi 1944 referred tentatively to Doleroleninae p. 401, Sdzuy.

Youngia uralica Tschernyschew 1893 p. 309 pl. lxxxiv figs. 4a-b, Lesnikova & Weber.

Yuepingia [sic] Lu 1956 see Niobinae subfam. n.

Yunnanocephalus Kobayashi 1936 referred tentatively to Bigotininae p. 401, Sdzuy.

Zacanthoides walapai sp. n. p. 194 pl. xx figs. 19–20 partes, Z. sp. undet. p. 194 pl. xix figs. 34–38 Cambrian (Middle) U.S.A. (Arizona), Resser Publ. Carneg. Instn No. 563 1945.

Zacanthoides sp. p. 84, Bright.

Zbirovia see Ectillaeninae subfam. n.

Zbirovia vaněki Šnajdr p. 48, Vaněk (3).

Zdicella see Ectillaeninae subfam. n.

Zelizskella [sic] deshayesi (Barr.), Z. sp. p. 47, Vaněk (3).

Zuninaspis see Thysanopyginae subfam. n.

Pygidia (resembling *Phillipsia scabra* Woodward) p. 177, **Caldwell.** 

Pygidium gen. and sp. undet. Cambrian (Upper) U.S.A. (Idaho) pl. lx fig. 22, Lochman & Hu.

Subgen. A see Calyptaulax.

Trilobite fragments p. 37ff., Thorsteinsson.

Unassigned pygidia nos. 9 and 10 of Rasetti 1945 belong to *Plethometopus 'possibly P. armatus*), Unassigned pygidium no. 7 of R. 1945 referred to *P. obtusus* p. 383, **Rasetti.** 

Undetermined pygidium No. 1 p. 118 pl. xxi fig. 7, Undetermined pygidium No. 2 p. 118 pl. xxi fig. 16 Ordovician (Lower), Berg & Ross.

## TRACE-FOSSILS POSSIBLY ATTRIBUTABLE TO TRILOBITES

Protichnites Owen 1852 discussed p. 7, P. sp. T. [? Cambrian or ? Ordovician] Australia (W. A.) p. 9 pls. figs. 2-7, Öpik (2).

Trilobite trails pl. v fig. b, pl. vii fig. a, McKee.

#### IV. \_ERRATUM

Z. R. 93 11 (for 1956).

The genera Acrocephalaspis, Kujandaspis, Kujandina, Ninaspis, Olentella, Pedinocephalus, Tatulaspis, Urbanaspis and their respective type species, although described as new in N. K. Ivshin's monograph of 1956, were first validly published by Ivshin in Izv. Acad. Sci. Kazakhstan S.S.R. [Geol.] 21 1955, and date therefrom.

## THE ZOOLOGICAL RECORD

THE object of The Zoological Record is to give, by means of an annual volume, complete lists of the works and publications relating to zoology in all its branches that have appeared during the year preceding the issue of the volume, together with a subject and systematic index.

The complete volume can be obtained from the Society at the price of £10 0s. 0d. per copy.

The following volumes are available:-

#### In Reprint Edition

y 9

.

.)

?

Vols. 1-30 (1864-1893). Obtainable from Messrs. Butterworths, 4-5, Bell Yard, Temple Bar, London, W.C.2.

Complete set of the first 25 volumes, bound in cloth, price £221 0s. 0d.

Complete set of 30 volumes, in paper covers, price £256 10s. 0d.

Separate volumes, in paper covers, vols. 1-20 £8 0s. 0d. each; vols. 21-25 £8 18s. 6d. each; vols. 26-30 £10 10s. 0d. each.

The "Insecta" section, obtainable separately, paper covers. Price complete set Vols. 1-30 £157 5s. 0d.; single volumes £5 5s. 0d. each.

#### In Original Edition

Vols. 65-71 (1928-1934) at £5 each. Vols. 86-91 (1948-1954) at £6 each. Vols. 92-95 (1955-1958) at £8 each.

For particulars of the availability and prices of separate Sections, application should be made to the Scientific Director, The Zoological Society of London, Regent's Park, London, N.W.1.

It should be noted that the separate Section Insecta can only be obtained from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

NOMENCLATOR ZOOLOGICUS.—Lists the names of all the genera and subgenera in zoology from the 10th Edition of Linnaeus 1758 to the end of 1945, with a bibliographical reference to the original description of each. Price 17½ guineas, postage extra. Price of separate volumes: Vols. 1-4 (1758-1935), Vol. 5 (1936-1945), 3½ guineas each, postage extra.

## SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The prices of the Sections from Volume 96 are:-

									S.	d.
1.	Comprehensive	e Zool	ogy		•••	•••	•••	•••	4	0
2.	Protozoa		•••	•••	•••	•••	•••		13	0
3.	Porifera	•••					•••		3	0
4.	Coelenterata				•••		•••	•••	5	0
5.	Echinodermata			•••					4	0
6.	Vermes								12	0
7.	Brachiopoda		•••			•••			4	0
8.	Bryozoa			•••					3	0
9.	Mollusca				•••	V.11.		•••	16	0
10.	Crustacea		•••			111			8	0
11.	Trilobita						•••		8	0
12.	Arachnida	•••			•••				14	0
13.	*Insecta				•••				60	0
14.	Protochordata	•••					•••		3	0
15.	Pisces		•••						11	0
16.	Amphibia		•••				•••		12	0
17.	Reptilia						•••		12	0
18.	Aves								12	0
19.	Mammalia		•••						12	0
20.	List of New G	Genera	and	Subgenera					4	0

\*Obtainable only from the Commonwealth Institute of Entomology, 56 Queen's Gate, London, S.W.7.

